

Topological boundary conditions, the BPS bound, and elimination of ambiguities in the quantum mass of solitons^{*}

Horatiu Nastase, Misha Stephanov, Peter van Nieuwenhuizen

Institute for Theoretical Physics,
S.U.N.Y. at Stony Brook, NY 11794-3840, USA

and

Anton Rebhan

Institut für Theoretische Physik, Technische Universität Wien,
Wiedner Hauptstr. 8-10, A-1040 Vienna, Austria

Abstract

We fix the long-standing ambiguity in the 1-loop contribution to the mass of a 1+1-dimensional supersymmetric soliton by adopting a set of boundary conditions which follow from the symmetries of the action and which depend only on the topology of the sector considered, and by invoking a physical principle that ought to hold generally in quantum field theories with a topological sector: for vanishing mass and other dimensionful constants, the vacuum energies in the trivial and topological sectors have to become equal. In the two-dimensional $N = 1$ supersymmetric case we find a result which for the supersymmetric sine-Gordon model agrees with the known exact solution of the S-matrix but seems to violate the BPS bound. We analyze the nontrivial relation between the quantum soliton mass and the quantum BPS bound and find a resolution. For $N = 2$ supersymmetric theories, there are no one-loop corrections to the soliton mass and to the central charge (and also no ambiguities) so that the BPS bound is always saturated. Beyond 1-loop there are no ambiguities in any theory, which we explicitly check by a 2-loop calculation in the sine-Gordon model.

^{*}The work is supported by NSF grant PHY97-22101.

1 Introduction

The calculation of quantum corrections to the mass of solitonic objects has been a subject of intense interest in the past [1] and has recently been revived in the light of a recent breakthrough in our understanding of non-perturbative dynamics in 3+1-dimensional supersymmetric (susy) gauge theories [2]. One of the most important ingredients of the non-perturbative analysis of such theories is the duality between extended solitonic objects, such as monopoles or dyons, and point-like particles. Another important ingredient is the concept of the BPS spectrum — the particles whose masses are proportional to their charges. Due to the supersymmetry algebra BPS states may be annihilated by the action of some of the supersymmetry generators and hence give rise to a smaller number of superpartners (“multiplet shortening”). Therefore the BPS value of the mass becomes a qualitative, rather than just a quantitative property [3].

Susy models in 1+1 dimensions provide a valuable nontrivial testing ground for these concepts. Solitons in these models are examples of BPS states whose mass at the classical level is proportional to the topological charge. The calculation of quantum corrections to the masses of these states has been subject to a long controversy. A number of one-loop calculations have been performed yielding different contradicting results [4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14].

The one-loop corrections to the mass of a soliton are given by

$$M_{sol}^{(1)} = \frac{1}{2} \sum (\omega^B - \omega^F) - \frac{1}{2} \sum (\tilde{\omega}^B - \tilde{\omega}^F) + \delta M \quad (1)$$

where $\omega^{B,F}$ are the energies of the small bosonic (fermionic) fluctuations about the classical soliton solution, $\tilde{\omega}$ are the corresponding energies of the linearized theory in the trivial vacuum, while δM is the counterterm which can be obtained from the expression for the classical mass of the soliton in terms of unrenormalized parameters by expanding into renormalized ones [15]. In order to make the above sums well defined, spatial boundaries are temporarily introduced to make the entire spectrum discrete.

One can identify two sources of ambiguities. As was discussed in [5], imposing different spatial boundary conditions on the small quantum fluctuations around the classical soliton gives different, sometimes even ultraviolet divergent, results. In this paper we present an analysis which answers the question: which boundary conditions are to be used in the one-loop calculation? The answer to this question can be found if one re-examines the original formulation of the problem. We consider the vacuum energy as a functional of the boundary conditions. We then single out a class of boundary conditions which do not introduce surface effects — the topological boundary conditions. They close the system on itself. There is a trivial (periodic) as well as a topologically non-trivial (with a Moebius-like twist) way of doing this. This definition of boundary conditions does not rely on a semiclassical loop expansion. We do not separate the classical part of the field from its quantum fluctuations; rather, the boundary conditions are imposed on the whole field. One then infers the correct boundary conditions for the classical part as well as for the quantum fluctuations from this single general condition.

Another source of ambiguity, as was pointed out recently [14], is the choice of the ultraviolet regularization scheme. The dependence on the choice of regularization scheme can

be understood as a peculiar property of those quantities, such as the soliton mass, which involve a comparison between two sectors with different boundary conditions, i.e., different topology. Indeed, the difference of the vacuum energies in the two sectors measures the mass of the soliton. The one-loop correction is then given by a sum over zero-point frequencies in the soliton sector which is quadratically divergent. A similar sum in another (the trivial) sector is to be subtracted in order to get an expression which is finite if written in terms of the renormalized parameters. It turns out that due to the bad ultraviolet behavior of both sums the result depends on the choice of the cut-off [14]. Cutting both sums off at the same energy in both sectors [7, 8, 9, 10] or taking equal numbers of modes in both sectors [16, 5, 12, 13] leads to different results. To add to the confusion, some authors do not include bound states and/or zero modes when they consider equal numbers of states in both sectors.¹ In this paper we propose a simple way of reducing the ultraviolet divergence of the sums over the zero-point energies, which eliminates the sensitivity to the ultraviolet cutoff. Instead of calculating the sums we calculate their derivative with respect to the physical mass scale in the theory. The constant of integration can be fixed by using the following observation: the vacuum energy should not depend on the topology when the mass is zero. This is the physical principle which allows us to perform the calculation unambiguously. It should be viewed as a renormalization condition.

From a practical point of view we need to use our condition that the vacuum energy functional does not depend on topology at zero mass only in the one-loop calculation. However, to preserve the spirit of our approach, we formulate this condition, as we do with our boundary conditions, for the full theory regardless of the semiclassical loop expansion. The mass that needs to be taken to zero is the physical, renormalized, mass scale. From this point of view this condition is a trivial consequence of dimensional analysis if we work with a renormalizable theory where all the physical masses are proportional to one mass scale. All the masses, including the soliton mass, vanish then at the same point, which is the conformal, or critical, point in the theory. Another way to look at our condition to fix the integration constant is to consider the Euclidean version of the 1+1 theory as a classical statistical field theory. Then the mass of the soliton is the interface tension between two phases. As is well known [17] the interface tension vanishes at the critical point; moreover, it vanishes with the same exponent as the inverse correlation length.

In section 2 we demonstrate our new unambiguous method of calculation using as an example the bosonic kink. We show that, as argued in [14], the correct result corresponds to mode number cutoff. (The same conclusion was recently reached for nontopological solitons in 3+1 dimensions [18].) In section 3 we apply our analysis of the topological boundary conditions to the case of an $N = 1$ susy soliton, where it leads to nontrivial consequences. We analyze the relation of our results to the BPS bound in section 4. In section 5 we analyze the $N = 2$ susy solitons and conclude that the one-loop corrections vanish completely. In section 6 we redo the 2-loop corrections for the case of the bosonic sine-Gordon soliton [19, 20], paying this time close attention to possible ambiguities, and find that no ultraviolet ambiguities appear. The ultraviolet ambiguity is thus purely a one-loop effect which leads to the interesting conjecture that it may be formulated in terms of a topological quantum

¹See for example the textbook [1], eq. (5.60). Actually in this reference the result of [16] is obtained but this requires neglecting a boundary term in the partial integration of (5.63).

anomaly.

2 Eliminating the one-loop ultraviolet ambiguity using a physical principle

In this section we present a general analysis regarding the calculation of the soliton mass which will help us eliminate the ultraviolet ambiguity discussed in [14]. We consider the ϕ^4 theory (kink) as an example, but the arguments can be applied to the sine-Gordon theory as well. The crucial property of these models from which our boundary conditions follow is the Z_2 symmetry $\phi \rightarrow -\phi$.

Let us take a step back from the actual calculation and try to define the mass of the soliton *before* we do the semiclassical expansion. We start from the observation that the soliton carries a conserved charge — the topological charge. This means that we can define the mass of the soliton as the difference between the energy of the system with nontrivial topology and the energy of the system with trivial topology. This definition coincides with the definition based on path integrals in the topological sector which are normalized by path integrals in the trivial sector [19]. The topological charge of the system is determined by the conditions at the spatial boundary. We view the vacuum energy as a functional of the boundary conditions. In general, a boundary condition could induce surface effects associated with the interaction of the system with the external forces responsible for the given boundary condition. We would like to avoid these contributions. There is a class of boundary conditions which do not produce such effects. These are what we call topological boundary conditions, which identify the degrees of freedom at different points on the boundary modulo a symmetry transformation. In our case there are two such possibilities: periodic and antiperiodic. These are dictated by the internal $Z(2)$ symmetry: $\phi(x) \rightarrow (-1)^p \phi(x)$, $p = 0, 1$. Crossing the boundary is associated with a change of variables leaving the action invariant: $\phi(-L/2) = (-1)^p \phi(L/2)$. The system behaves continuously across the boundary, only our description changes. In effect such boundary conditions do not introduce a boundary, rather they close a system in a way similar to the Moebius strip.²

We spent so much time on this, perhaps, trivial point in order to make the choice of boundary conditions for the theory with fermions clear. The analysis of fermions, however, will be postponed until the next section. In the literature a large number of other boundary conditions have been considered, both in the bosonic and in the fermionic sectors but from our perspective they all introduce surface effects or are even inconsistent.

It should now be clear that the mass of the soliton can be defined as the difference of the vacuum energy with antiperiodic and with periodic boundary conditions when the volume $L \rightarrow \infty$. This definition does not rely on the semiclassical expansion. Returning now to the semiclassical calculation we see that at the classical level the equations of motion select the trivial or the soliton vacuum configuration depending on the topology. Less trivially, we see that at the one-loop level the boundary conditions that should be used for the small fluctuations about the soliton configuration should be antiperiodic. We must point out that

²Though topological boundary conditions do not induce boundary effects, finite volume effects vanishing as $L \rightarrow \infty$ are of course present. Such effects will be discussed in section 4.

the choice of boundary conditions does not affect the result of the calculation in the purely bosonic case. We shall nevertheless use antiperiodic boundary conditions in the soliton sector in this section to be faithful to our nonperturbative definition of the soliton mass. We shall see in the next section that for fermions the choice of the boundary conditions becomes crucial.

A few points about the classical antiperiodic soliton should be stressed here. The topological boundary condition reads

$$\phi(-L/2) = (-1)^p \phi(L/2) \quad \text{and} \quad \phi'(-L/2) = (-1)^p \phi'(L/2), \quad (2)$$

where the nontrivial sector is selected when $p = 1$. Note that the derivative with respect to x , ϕ' , must also be antiperiodic in the soliton sector. The classical soliton solution $\phi(x)$ can be viewed as a trajectory of a particle with coordinate ϕ moving in time x in the potential $-V(\phi)$. The particle is oscillating about the origin $\phi = 0$ with a period which depends on the amplitude. When the period is equal to $2L$ the trajectory during half of the period is the antiperiodic soliton satisfying the boundary conditions (2). The endpoints of the trajectory need not necessarily be the turning points. For example, the particle at time $-L/2$ can start downhill at some ϕ_0 with nonzero velocity, then pass the point at the same height on the opposite side, i.e., $-\phi_0$, going uphill, then turn and after that at time $L/2$ pass the point $-\phi_0$ again, but going downhill. Clearly, for this trajectory, (2) is satisfied, whereas restricting the usual soliton solution centered at $x = 0$ to the interval $(-L/2, L/2)$ would lead to a solution for which $\phi(-L/2) = -\phi(L/2)$, but $\phi'(-L/2) = +\phi'(L/2)$. When $L \rightarrow \infty$ the turning points of the trajectory come infinitesimally close to the minima of $V(\phi)$ and we recover the usual $L = \infty$ soliton.

Next, we want to address the problem of the ultraviolet ambiguity in the one-loop calculation. To summarize the beginning of this section

$$M \equiv E_1 - E_0, \quad (3)$$

where E_p , $p = 0, 1$ are the energies of the system with the periodic and antiperiodic boundary conditions of (2) respectively. At the classical level this gives $M_{\text{cl}} \sim m^3/(3\lambda)$ for the kink, where m is the mass of the elementary boson at tree level, and λ the dimensionful coupling constant. The order $\hbar$ correction is due to the fact that boundary conditions change the spectrum of zero-point fluctuations. Two factors are responsible for the ultraviolet ambiguity discussed in [14]. One is the fact that the terms in the sums over zero-point energies grow making each sum strongly (quadratically) ultraviolet cutoff dependent. Second is that one has to compare the spectrum in two different vacua. Taking all the modes below a certain energy in both systems leads to a different result than taking equal numbers of modes. It would be nice if there was a parameter in the theory whose variation would continuously interpolate between the two vacua. This is not possible due to the topological nature of the difference between the vacua. However, we can identify a certain value of the dimensionful parameters of the theory at which the vacuum energy should become independent of the topology. This will be one ingredient of our calculation.

Another ingredient is the observation that one can reduce the ultraviolet divergence of the sums of zero-point energies by differentiating w.r.t. m . The terms in the differentiated

sums become then decreasing and as a result the sums (now only logarithmically divergent) can be unambiguously calculated. But the price is that we need to supply the value for the integration constant to recover the function from its derivative. This can be done using a physical principle that relates the energies of the two vacua at some value of the mass. One must realize that the difference in the energies arises because of the nontrivial potential for the scalar field. If this potential vanishes the energies of the two vacua become equal. In the absence of the potential the mass m of the boson is zero and the soliton disappears. Therefore the constant of integration over m is fixed by the condition that the energy difference between the two vacua must vanish when $m \rightarrow 0$. A subtlety here is that m should be sufficiently large compared to $1/L$ so that finite volume effects can be neglected. The limit $m \rightarrow 0$ should be understood in the sense that the mass approaches $O(1/L)$, where L is large. Then the difference between the vacuum energies must be $O(1/L)$. Also note, that other dimensionful parameters in the theory should be scaled accordingly when $m \rightarrow 0$, e.g., $\lambda/m^2 = \text{const}$ in the $\lambda\phi^4$ theory.

We want to relate the mass of the soliton to other parameters of the theory. The relation to the bare parameters m_0 and λ will contain infinities. The infinities in the relation of physical quantities to the bare parameters in this theory should be eliminated if we renormalize the mass $m_0 = m + \delta m$, where

$$\delta m = \frac{3\lambda}{2m} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{\sqrt{k^2 + m^2}}. \quad (4)$$

With this renormalization of m_0 tadpole diagrams vanish. In the ϕ^4 theory the physical pole mass of the meson differs from m by a *finite* amount $-\sqrt{3}\lambda/(4m)$ [14]; however, it suffices to use m for our purposes. If we rerun this analysis for the sine-Gordon theory, the tadpole renormalized mass m would, at one-loop order, coincide with the physical meson mass. If we use this renormalized mass in the expression for the soliton mass we get an additional one-loop counterterm, δM ,

$$\frac{m_0^3}{3\lambda} = \frac{m^3}{3\lambda} + \delta M, \quad (5)$$

where

$$\delta M = \frac{m^2 \delta m}{\lambda} = \frac{3m}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{\sqrt{k^2 + m^2}}. \quad (6)$$

Now we differentiate the well-known expression for the one-loop correction $M^{(1)}$ to the soliton mass with respect to the mass m

$$\frac{dM^{(1)}}{dm} = \frac{d(\delta M)}{dm} + \frac{1}{2} \sum_n \frac{d\omega_n}{dm} - \frac{1}{2} \sum_n \frac{d\tilde{\omega}_n}{dm}. \quad (7)$$

For the spectrum $\tilde{\omega}_n$ in the trivial sector one obtains

$$\frac{d\tilde{\omega}_n}{dm} = \frac{m}{\sqrt{\tilde{k}_n^2 + m^2}}, \quad \tilde{k}_n L = 2\pi n. \quad (8)$$

For the soliton sector

$$\frac{d\omega_n}{dm} = \frac{1}{\sqrt{k_n^2 + m^2}} \left(m + k_n \frac{dk_n}{dm} \right) = \frac{1}{\sqrt{k_n^2 + m^2}} \left(m + \frac{1}{L} \frac{k_n^2}{m} \delta'(k_n) \right),$$

$$k_n L + \delta(k_n) = 2\pi n + \pi, \quad (9)$$

where we used the fact that $\delta(k)$ depends on m only through k/m to convert the derivative w.r.t. m into the derivative w.r.t. k .

We convert the sums over the spectrum in (7) into integrals over k using

$$\sum_n f(\tilde{k}_n) = L \int_{-\infty}^{\infty} \frac{dk}{2\pi} f(k) + O(1/L); \quad (10)$$

and

$$\sum_n f(k_n) = L \int_{-\infty}^{\infty} \frac{dk}{2\pi} f(k) \left(1 + \frac{\delta'(k)}{L} \right) + O(1/L). \quad (11)$$

These expressions follow from the Euler-Maclaurin formula which is valid for a smooth function $f(k)$ vanishing at $k = \infty$. In our case $f(\tilde{k}) = d\tilde{\omega}/dm$ and $f(k) = d\omega/dm$ satisfy these conditions. From the Euler-Maclaurin formula one can also see that in the naive calculation with $f(k) = \omega$ the ambiguous contribution, which comes from regions δ/L at the ultraviolet ends of the integration interval, is non-vanishing due to the fact that $f(k)$ grows with k .




Figure 1: The left- and the right-hand sides of the equation $\delta(k) = 2\pi n + \pi - kL$ are plotted schematically by solid lines in the case of the ϕ^4 kink (two bound states). The dashed line represents the value of $\delta(k)$ without the discontinuity $2\pi\varepsilon(k)$. Observe that with this discontinuity the mode numbers $n = -1, 0$ should be left out, while the spectrum of allowed values of k is not affected.

We can use the following expression for the phase shifts $\delta(k)$ in the case of the kink in ϕ^4 theory:

$$\delta(k) = \left(2\pi - 2 \arctan \frac{3m|k|}{m^2 - 2k^2} \right) \varepsilon(k), \quad (12)$$

where we added the term $2\pi\varepsilon(k)$ to ensure that $\delta(|k| \rightarrow \infty) \rightarrow 0$. As a result, $\delta(k)$ is discontinuous at $k = 0$: $\delta(0_{\pm}) = \pm 2\pi$. It is then easy to see (Fig. 1) that for $n = -1$ and $n = 0$ the equation (9) does not have solutions for k . It is pleasing to observe that this defect is matched by the existence of two discrete modes: $\omega_0 = 0$ (the translational zero mode) and $\omega_{-1} = \sqrt{3}m/2$ (a genuine bound state). To these discrete modes we can assign (somewhat arbitrarily) those “unclaimed” n ’s. That this matching is not a coincidence follows from Levinson’s theorem: $\delta(0_{\pm}) = \pm\pi n_{\text{ds}}$, where n_{ds} is the number of discrete solutions. Since the discontinuity in $\delta(k)$ is an integer multiple of 2π , it does not change the spectrum of the allowed values of k (Fig. 1). This spectrum near the origin is given by $kL = \dots, -3\pi, -\pi, \pi, 3\pi, \dots$ up to $O(1/L)$ for δ either with or without the $2\pi\varepsilon(k)$ term in (12), and the values of $\delta'(k)$ and $f(k)$ on this set of k ’s are also not affected.

Putting now all the pieces together we obtain

$$\frac{dM^{(1)}}{dm} = \frac{d(\delta M)}{dm} + \frac{1}{2} \sum_{\text{ds}} \frac{d\omega_{\text{ds}}}{dm} + \frac{1}{2m} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \sqrt{k^2 + m^2} \delta'(k). \quad (13)$$

This formula is universal. Substituting for the ϕ^4 theory the particular values for δM from (6) and $\delta(k)$ from (12) we find³

$$\begin{aligned} \frac{dM^{(1)}}{dm} &= -\frac{3m^2}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{(k^2 + m^2)^{3/2}} + \frac{\sqrt{3}}{4} \\ &\quad - \frac{3m^2}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{\sqrt{k^2 + m^2}(m^2 + 4k^2)} = \frac{1}{4\sqrt{3}} - \frac{3}{2\pi}. \end{aligned} \quad (14)$$

Integrating over m and using that $M^{(1)} = 0$ when $m = 0$ we obtain the result for the one-loop correction to the kink mass which was previously obtained using mode number cutoff [16, 14]

$$M^{(1)} = m \left(\frac{1}{4\sqrt{3}} - \frac{3}{2\pi} \right). \quad (15)$$

3 Fermions, supersymmetry, and topological boundary conditions

In this section we shall extend the ideas introduced in the previous section to theories with fermions, and in particular theories with supersymmetry. The following analysis can be applied to any $N = (1, 1)$ supersymmetric theory with Lagrangian

$$\mathcal{L} = -\frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - \frac{1}{2} U^2(\phi) - \frac{1}{2} (\bar{\psi} \not{\partial} \psi + U'(\phi) \bar{\psi} \psi), \quad (16)$$

where $U(\phi)$ is a symmetric function, admitting a classical soliton solution. For example, for the kink $U(\phi) = \sqrt{\lambda/2}(\phi^2 - \frac{m_0^2}{2\lambda})$. We use $\{\gamma^{\mu}, \gamma^{\nu}\} = 2g^{\mu\nu}$ with $g^{00} = -1$, $g^{11} = 1$, and ψ is

³Use $\int_{-\infty}^{\infty} (1+x^2)^{-1/2} (1+4x^2)^{-1} dx = 2\sqrt{3}\pi/9$ and $\int_{-\infty}^{\infty} (1+x^2)^{-3/2} dx = 2$.

a Majorana spinor: $\bar{\psi} = \psi^\dagger i\gamma^0 = \psi^T C$ with $C\gamma^\mu = -(\gamma^\mu)^T C$. The action is invariant under $\delta\phi = \bar{\epsilon}\psi$ and $\delta\psi = (\not{\partial}\phi - U)\epsilon$.

First of all we want to identify the class of topological boundary conditions. The discrete transformation taking $\phi \rightarrow -\phi$ must be accompanied by $\psi \rightarrow \gamma_3\psi$ with $\gamma_3 = \gamma^0\gamma^1$ to leave the action invariant. From this symmetry transformation we obtain topological boundary conditions

$$\begin{aligned}\phi(-L/2) &= (-1)^p \phi(L/2), & \phi'(-L/2) &= (-1)^p \phi'(L/2), \\ \psi(-L/2) &= (\gamma_3)^p \psi(L/2), & p &= 0, 1.\end{aligned}\tag{17}$$

The value $p = 0$ gives a trivial periodic vacuum while $p = 1$ selects a nontrivial soliton vacuum.

As one can see, the reasons behind our choice of the topological boundary conditions do not include supersymmetry. The same arguments apply to any theory with a Yukawa-like interaction between fermions and bosons. From this point of view it is very gratifying to discover that the $p = 1$ topological boundary condition (17) preserves half of the supersymmetry of the Lagrangian (16). An easy way to see that is to consider the Noether current corresponding to the supersymmetry

$$J^\mu = -(\not{\partial}\phi + U)\gamma^\mu\psi.\tag{18}$$

Integrating the conservation equation $\partial_\mu J^\mu = 0$ over space we find

$$\frac{\partial Q}{\partial t} \equiv \frac{\partial}{\partial t} \int_{-L/2}^{L/2} dx J^0(x) = -[J^1(x)]_{-L/2}^{L/2},\tag{19}$$

where the r.h.s. is simply the total current flowing into the system. Using the boundary condition (17) we obtain

$$-[J^1(x)]_{-L/2}^{L/2} = [(\not{\partial}\phi + U)\gamma^1\psi - (-\not{\partial}\phi + U)\gamma^1\gamma^3\psi]_{L/2} = (1 + \gamma_3)(\not{\partial}\phi + U)\gamma^1\psi|_{L/2}.\tag{20}$$

We see that the $(1 - \gamma_3)$ projection of the supercharge Q is conserved. Note that different projections of Q are *classically* conserved on the soliton or the antisoliton background. The soliton with $\phi' + U = 0$ and $\psi = 0$ is invariant under a susy transformation with a parameter ϵ if $(1 + \gamma^1)\epsilon = 0$. This means the projection P_+Q (with $P_\pm = 1 \pm \gamma_1$) of the supercharge vanishes on the soliton configuration to linear order in the quantum fields. For the antisoliton P_-Q has this property. This should be expected since the topological boundary condition does not distinguish between the soliton and the antisoliton.

Similarly one can see that the topological boundary condition does not break translational invariance. The conservation equation for the stress tensor reads $\partial_\mu T^{\mu\nu} = 0$, where

$$T^{\mu\nu} = \partial^\mu\phi\partial^\nu\phi + \frac{1}{4}(\bar{\psi}\gamma^\mu\partial^\nu\psi + \bar{\psi}\gamma^\nu\partial^\mu\psi) + \mathcal{L}g^{\mu\nu}.\tag{21}$$

In general, the non-conservation of total momentum is again due to the boundary term

$$\frac{\partial P}{\partial t} \equiv \frac{\partial}{\partial t} \int_{-L/2}^{L/2} T^{01} = -[T^{11}]_{-L/2}^{L/2}.\tag{22}$$

We see that the defining property of the topological boundary condition, that it relates the fields at $-L/2$ and $L/2$ by a transformation leaving $\mathcal{L}$ invariant, ensures momentum conservation.

Note also that there is another $Z(2)$ symmetry in the Lagrangian (16): $\psi \rightarrow (-1)^q \psi$. This can be used to extend the set of topological boundary conditions (17) to

$$\begin{aligned}\phi(-L/2) &= (-1)^p \phi(L/2), & \phi'(-L/2) &= (-1)^p \phi'(L/2), \\ \psi(-L/2) &= (-1)^q (\gamma_3)^p \psi(L/2), & p, q &= 0, 1.\end{aligned}\quad (23)$$

The values $(p, q) = (0, 0)$ give a topologically trivial sector. The sector $(0, 1)$ is also trivial, but the fermions have a twist. Two classically nontrivial vacua are obtained with $p = 1$ and $q = 0, 1$. For $p = 1$ the two values of q correspond to the arbitrariness of the sign choice of the γ_3 matrix, and are related to each other by space parity transformation $\psi(x, t) \rightarrow \gamma^0 \psi(-x, t)$. Therefore one should expect $E(1, 0) = E(1, 1)$, which one can check is true at one-loop.

As in the previous section we define the mass of the soliton as the difference of the energies E_p of these vacua: $M \equiv E_1 - E_0$. At the classical level one finds $M = M_{\text{cl}}$, where M_{cl} is the classical soliton mass. The one-loop correction is determined by integrating

$$\frac{dM^{(1)}}{dm} = \frac{d(\delta M)}{dm} + \frac{1}{2} \sum_n \frac{d\omega_n^B}{dm} - \frac{1}{2} \sum_n \frac{d\omega_n^F}{dm} - \left(\frac{1}{2} \sum_n \frac{d\tilde{\omega}_n^B}{dm} - \frac{1}{2} \sum_n \frac{d\tilde{\omega}_n^F}{dm} \right) \quad (24)$$

over m . The expressions for the derivatives $d\omega_n^B/dm$ and $d\tilde{\omega}_n^B/dm$ are the same as in the bosonic case, see (8) and (9). In order to find the corresponding expressions for the fermionic frequencies we need to obtain the quantization condition for k_n . For the trivial sector we have simply $\tilde{k}_n L = 2\pi n$.

The nontrivial sector requires more careful analysis. The frequencies ω are obtained by finding solutions of the equation

$$(\gamma^\mu \partial_\mu + U') \psi = 0 \quad (25)$$

of the form $\psi = \psi(x) \exp\{-i\omega t\}$. Multiplying this equation by $(-\gamma^\mu \partial_\mu + U')$ we find

$$\left(-\partial^2 + (U')^2 + \gamma^1 U'' \right) \psi = 0, \quad (26)$$

where we used $\phi'_{\text{sol}} = -U(\phi_{\text{sol}})$, which follows from the classical equation of motion for the soliton in the $L \rightarrow \infty$ limit. Projecting this equation using $P_\pm = (1 \pm \gamma^1)/2$ we see that ψ_+ (where $\psi_\pm \equiv P_\pm \psi$) obeys the same equation as the bosonic small fluctuations, hence

$$\psi_+ = \alpha_+ e^{-i\omega t + i(kx \pm \delta/2)} \quad \text{when } x \rightarrow \pm\infty. \quad (27)$$

The ψ_- component can then be obtained by acting with P_+ on (25)

$$\partial_0 \gamma^0 \psi_- + (\partial_1 + U') \psi_+ = 0. \quad (28)$$

which together with (27) gives the asymptotics of ψ_-

$$\psi_- = -\gamma^0 \alpha_+ \frac{k \mp im}{\sqrt{k^2 + m^2}} e^{-i\omega t + i(kx \pm \delta/2)} = -\gamma^0 \alpha_+ e^{\pm i\theta/2} e^{-i\omega t + i(kx \pm \delta/2)} \quad \text{when } x \rightarrow \pm\infty, \quad (29)$$

where $\theta(k) = -2 \arctan(m/k)$ and we used the fact that in the $L \rightarrow \infty$ limit $U' \rightarrow \pm m$. Therefore the solutions of (25) have asymptotics

$$\psi = \psi_+ + \psi_- = \left(1 - \gamma^0 e^{\pm i\theta/2}\right) \alpha_+ e^{-i\omega t + i(kx \pm \delta/2)} \quad \text{when } x \rightarrow \pm\infty, \quad (30)$$

where α_+ is the eigenvector of γ^1 with eigenvalue $+1$.

Although one could continue the derivation without specifying the representation for the γ matrices (an exercise for the reader) we find it more concise to adopt a certain representation. The most convenient is the following one in terms of the Pauli matrices: $\gamma^0 = -i\tau_2$, $\gamma^1 = \tau_3$, and hence $\gamma_3 = \tau_1$. It has two advantages. First, α_+ has now only an upper component. Second, the Majorana condition becomes simply $\psi^* = \psi$ and the equation (25) is real. In this representation (30) becomes

$$\psi = \begin{pmatrix} 1 \\ -e^{\pm i\theta/2} \end{pmatrix} \alpha e^{-i\omega t + i(kx \pm \delta/2)} \quad \text{when } x \rightarrow \pm\infty, \quad (31)$$

where α is a complex number.

Now we impose the boundary condition $\psi(-L/2) = \gamma_3 \psi(L/2)$. The field ψ in equation (25) must be real. This means that only the real part of (31) need to satisfy the boundary condition. This condition should, however, be maintained for all t . Therefore, due to the oscillating phase $\exp(-i\omega t)$, a complex equation must be satisfied

$$e^{-i(kL+\delta)} \begin{pmatrix} 1 \\ -e^{-i\theta/2} \end{pmatrix} = \Gamma \begin{pmatrix} 1 \\ -e^{i\theta/2} \end{pmatrix} \quad (32)$$

We introduced $\Gamma = \gamma_3$ in order to discuss briefly the following point. One could consider a more general boundary condition: $\psi(-L/2) = \Gamma \psi(L/2)$ with some matrix Γ (real in the Majorana representation we have chosen). One can see from (32) that certain boundary conditions cannot be satisfied, for example, the frequently employed [7, 10, 11, 12, 13, 14] periodic boundary conditions with $\Gamma = 1$. Equation (32) provides an additional consistency check for our choice of boundary condition.

With the topological boundary condition $\Gamma = \gamma_3 = \tau_1$, we find that equation (32) is satisfied provided

$$kL + \delta + \frac{\theta}{2} = 2\pi n + \pi. \quad (33)$$

Using this quantization rule we find

$$\frac{d\omega_n^F}{dm} = \frac{1}{\sqrt{k^2 + m^2}} \left(m + k_n \frac{dk_n}{dm} \right) = \frac{1}{\sqrt{k^2 + m^2}} \left(m + \frac{1}{L} \frac{k_n^2}{m} \left(\delta'(k) + \frac{\theta'(k)}{2} \right) \right) \quad (34)$$

Now we convert the sums over modes into integrals. For the bosonic and fermionic sums in the trivial sector formula (10) applies and the sums cancel each other (no cosmological constant in the trivial susy vacuum). For the bosonic sum in the nontrivial sector we use again (11). For the fermionic sum a formula analogous to (11) applies with $\delta + \theta/2$ instead of δ , which follows from (33). Again, as for the bosonic modes, due to the discontinuity in δ there are n_{ds} values of n which do not lead to a solution of (33). The remaining n lead to k




Figure 2: The left- and the right-hand sides of the equation $\delta + \theta/2 = 2\pi n + \pi - kL$ are plotted schematically in the case of the supersymmetric ϕ^4 kink (two bound states). The dashed line represent the value of $\delta(k) + \theta(k)/2$ without the discontinuity $2\pi\epsilon(k)$. As in the bosonic spectrum the discontinuity leads to $n = -1, 0$ mode numbers being skipped, while the spectrum of allowed values of k is not affected.

values (namely, in the case of the kink: $kL = \dots, -5\pi/2, -\pi/2, 3\pi/2, 7\pi/2, \dots$, see Fig. 2) which in the limit $L \rightarrow \infty$ give a continuous integration measure $Ldk/(2\pi)$ near $k = 0$.

Putting now everything into formula (24) we find

$$\frac{dM^{(1)}}{dm} = \frac{d(\delta M)}{dm} - \frac{1}{4m} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \sqrt{k^2 + m^2} \theta'(k). \quad (35)$$

The one-loop mass counterterm is given by [14]

$$\delta M = \frac{m}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{\sqrt{k^2 + m^2}}. \quad (36)$$

It follows from the renormalization counterterm δm which is chosen to cancel the sum of the bosonic and fermionic tadpole diagrams. Substituting into (35) we find $dM^{(1)}/dm = -1/(2\pi)$, hence

$$M^{(1)} = -\frac{m}{2\pi}. \quad (37)$$

This result differs from the one two of us have obtained previously [14] using a mode-number regularization scheme with the conventionally employed (but, as we have argued, untenable) periodic boundary conditions.

In the special case of the supersymmetric sine-Gordon model, we can compare this result with the one obtained from the Yang-Baxter equation assuming the factorization of the S-matrix [21]. The mass spectrum is then given by [22]

$$m_n = 2M \sin(n\gamma/16), \quad (38)$$

where γ in the notation of ref. [22] is related to the bare coupling β through

$$\frac{1}{\gamma} = \frac{1 - \beta^2/4\pi}{4\beta^2} = \frac{1}{4\beta^2} - \frac{1}{16\pi}. \quad (39)$$

Expanding (38) for $n = 1$ we find

$$m_1 = \frac{M\gamma}{8} + O(\gamma^3). \quad (40)$$

Since this is the lightest mass in the spectrum we identify it with the meson mass $m = m_1$. Taking the ratio M/m_1 and using (39) we obtain

$$\frac{M}{m} = \frac{8}{\gamma} + O(\gamma) = \frac{2}{\beta^2} - \frac{1}{2\pi} + O(\beta^2). \quad (41)$$

The first term is the classical result, the second is the 1-loop correction. This means that the 1-loop correction to M following from the exact S-matrix factorization calculation [22] is the same as our 1-loop result (37).

The next question we address is whether such a negative correction is in agreement with the well-known BPS bound. This question has been subject to controversy and deserves a separate section.

4 Quantum BPS bound, soliton mass, and finite size effects

As was first realized by Olive and Witten [3], the naive supersymmetry algebra in a topologically nontrivial sector is modified by central charges. The susy generators for the $N = 1$ model read in the representation of section 3

$$Q_{\pm} \equiv P_{\pm} \int_{-L/2}^{L/2} [-(\not{\partial}\varphi + U)\gamma^0\psi]dx = \int_{-L/2}^{L/2} [\dot{\varphi}\psi_{\pm} + (\varphi' \pm U)\psi_{\mp}]dx, \quad (42)$$

where $P_{\pm}$ is again $(1 \pm \gamma_1)/2$. Using canonical commutation relations we arrive at the following algebra

$$\{Q_{\pm}, Q_{\pm}\} = 2H \pm 2Z; \quad \{Q_+, Q_-\} = 2P, \quad (43)$$

where

$$H \equiv \int_{-L/2}^{L/2} \left[\frac{1}{2}\dot{\varphi}^2 + \frac{1}{2}(\varphi')^2 + \frac{1}{2}U^2 + \frac{i}{2}(\psi_+\psi'_- + \psi_-\psi'_+) - iU'\psi_+\psi_- \right] dx \quad (44)$$

$$P \equiv \int_{-L/2}^{L/2} \left[\dot{\varphi}\varphi' + \frac{i}{2}(\psi_+\psi'_+ + \psi_-\psi'_-) \right] dx \quad (45)$$

$$Z \equiv \int_{-L/2}^{L/2} \varphi' U dx \quad (46)$$

The central charge Z is clearly a boundary term.

Let us comment on some subtleties in the derivation of (43). The Dirac delta functions in the equal-time canonical commutation relations can be written as $\delta(x, y) = \sum_m \eta_m(x) \eta_m(y)$,

where η_m is a complete set of functions *satisfying the boundary conditions* of the corresponding field. For such $\delta(x, y)$ one has:

$$\int_{-L/2}^{L/2} \phi(x) \delta(x - L/2) = \frac{\phi(L/2) \pm \phi(-L/2)}{2} \quad (47)$$

$$\int_{-L/2}^{L/2} \phi(x) \delta(x + L/2) = \frac{\phi(-L/2) \pm \phi(L/2)}{2} \quad (48)$$

For the bosons in the topological sector we need the $-$ signs. For the fermions $\psi_+(x) + \psi_-(x)$ one needs the $+$ signs, but for the fermions $\psi_+(x) - \psi_-(x)$ one needs the $-$ signs. That *some* subtlety in the delta functions is present, is immediately clear if one considers the double integral $\int \int dx dy f(x) \partial_x \delta(x - y) g(y)$, and either directly partially integrates the derivative $\partial/\partial x$, or first replaces $\partial_x \delta(x - y)$ by $-\partial_y \delta(x - y)$ and then partially integrates w.r.t. y . One gets the same result provided

$$f g(x)|_{x \in B} = \int_{-L/2}^{L/2} dy f(x) \delta(x - y) g(y)|_{x \in B} + \int_{-L/2}^{L/2} dx f(x) \delta(x - y) g(y)|_{y \in B} \quad (49)$$

where the notation $h(x)|_{x \in B}$ implies $h(L/2) - h(-L/2)$. Naively, there is a factor of 2 missing in this equation, but with the more careful definitions of $\delta(x - y)$ for periodic or antiperiodic functions, consistency is obtained.

With these delta functions one finds that the boundary terms in the $\{Q_\pm, Q_\pm\}$ anticommutators reproduce the consistency condition (49), hence cancel, whereas in the $\{Q_+, Q_-\}$ anticommutator one finds the boundary term

$$-i \int dx \psi_+(x) \delta_a(x - y) \psi_+(y)|_{y \in B} - i \int dx \psi_-(x) \delta_a(x - y) \psi_-(y)|_{y \in B} \quad (50)$$

where $\delta_a(x - y)$ is the bosonic (antisymmetric) delta function defined above. These terms cancel if one uses our topological boundary conditions. In the $\{Q, Q\}$ relations one does not encounter subtleties involving delta functions for the fermions because there are no derivatives of fermions in $Q_\pm$.

Since the operators $Q_\pm$ are hermitian, one finds that the following relation exists between the expectation values of operators H and Z :

$$\langle s | H | s \rangle \geq |\langle s | Z | s \rangle|, \quad (51)$$

for any quantum state s .

As we have already pointed out in section 3 (20), only one linear combination of Q_+ and Q_- is conserved in the soliton sector: $Q_+ \pm Q_-$ for $q = 1, 0$ respectively. Taking for definiteness $q = 0$ we derive from (42), (44) the following commutation relations:

$$i[H, Q_+ + Q_-] = 2[(\dot{\varphi} + \varphi')(\psi_+ + \psi_-) + U(\psi_+ - \psi_-)]_{x=L/2} \quad (52)$$

$$i[P, Q_+ + Q_-] = 2[(\dot{\varphi} + \varphi')(\psi_+ + \psi_-) - U(\psi_+ - \psi_-)]_{x=L/2}, \quad (53)$$

while the other linear combination $Q_+ - Q_-$ commutes with both H and P .⁴ We also find that the operator Z does not commute with the Hamiltonian in the soliton sector:

$$i[H, Z] = 2U\dot{\varphi}\Big|_{x=L/2} \quad (54)$$

⁴In these relations one must use the proper definitions of the delta functions for the fermions since derivatives of $\psi_\pm$ appear in H . As one can see, (52) agrees with (20).

Let us examine carefully the meaning of this last result. Strictly speaking, it implies that Z is not a central charge. We shall show that this fact reflects a certain property of Z in a *finite* volume L . Let us ask the following question: what is the expectation value of Z in the soliton vacuum state $|\text{sol}\rangle$? It is clear that for any finite volume L : $\langle \text{sol} | Z | \text{sol} \rangle = 0$, since neither a positive nor a negative value is distinguished by our boundary conditions. This is a consequence of the fact that H and Z do not commute. The ground state $|\text{sol}\rangle$ is an eigenvector of H , but it need not be and is not an eigenvector of Z .

To make the next step clear it is convenient to use the following observation: The value of Z measures the position of the soliton. Indeed, for the classical configuration if the center of the soliton is at $x = 0$ exactly then Z is M (up to $O(1/L)$). If we now move the center of the classical soliton the value of Z will decrease and reach zero when the center of the soliton is at $L/2$. If we continue shifting the solution in the same direction, and bearing in mind its antiperiodicity, Z will become increasingly negative and it will reach the value $-M$ when there is an antisoliton at $x = 0$. If we deal with a configuration which is a distortion of the soliton, the center of the soliton is not well defined but Z is and can give one an idea of where the soliton is. On the quantum level this corresponds to the fact that P and Z do not commute. If we act with P (this is our spatial shift) on an eigenstate of Z we generate a different eigenstate of Z and the expectation value of Z changes.

Now, if we think of Z as (a nonlinear function of) the coordinate and P as the momentum of the soliton, the next step becomes clear. The vacuum is an eigenstate of P with eigenvalue 0. Therefore, it is a superposition of the eigenstates of Z . Positive and negative values of Z enter with the same weight into this superposition (the corresponding eigenvectors can be obtained from each other by acting with $\exp(iLP)$ — a shift by L). Therefore $\langle \text{sol} | Z | \text{sol} \rangle = 0$ for any finite L .

It is, however, too early to conclude that the equation (51) does not lead to any condition on $\langle \text{sol} | H | \text{sol} \rangle$ apart from semipositivity. The expectation value of Z can be compared to an order parameter in a system with spontaneous symmetry breaking. It is only nonzero if the thermodynamic limit $L \rightarrow \infty$ is taken properly. We shall now analyze how the limit $L \rightarrow \infty$ must be taken in the case of the soliton.

One can view the soliton as an almost classical particle (as long as the coupling constant is small) which is subject to Brownian motion due to quantum fluctuations. This is the meaning of the fact that H and Z do not commute: Z , or the position of the soliton, depends on time. If we wait for a sufficiently long time it will cover all possible positions and the expectation value of Z will be zero. However, it is obvious that for large L most of the time the soliton will spend away from the boundaries. It means that if one starts from a state of the soliton away from the boundary so that $Z = M$ and limits the time of observation the expectation value of Z will remain close to M .

How long does it take for the soliton to cover all the volume L ? Since it is a random walk the distance from the original position is $O(\sqrt{t})$ and it will stay away from the boundary if we restrict the interval t of the observation to $t \ll O(L^2)$. Such a restriction will mean that we introduce an error of at most $O(1/L^2)$ in the energy due to uncertainty principle. This is small when L is large. Alternatively, one can do Euclidean rotation and consider classical statistical theory of an interface in 2 dimensions. The random walk in this case is the well-known roughening of the interface [17]. It leads to smearing of the interface to the

width of $w = \text{const} \sqrt{t/M}$, where M is the one-dimensional interface tension, which is the soliton mass. The correction to the tension turns out to be $O(1/t)$, as we have already seen. (More rigorously, the partition function of the wall is not just $\exp(-Mt)$ in the $t \rightarrow \infty$ limit but has a preexponent $L\sqrt{M/t}$ due to the fluctuations of the interface. The factor L arises from the integration over the volume of the collective coordinate with a familiar measure $\sqrt{Mt}$, and an additional factor $1/t$ comes from the determinant of nonzero soft vibrational modes of the interface [23]).

Therefore the thermodynamic limit $L \rightarrow \infty$ in our system which gives nonzero $\langle \text{sol}|Z|\text{sol} \rangle$ corresponds to the energy measurement whose duration t is small compared to ML^2 . The bound (51) must apply to the result of such a measurement. The expectation value of Z in such a thermodynamic limit is what was calculated in [8, 14]:

$$\begin{aligned} Z &= Z_{\text{cl}}(m_0) + \int_{-\infty}^{\infty} dx \frac{d}{dx} \left[\frac{1}{2} U'(\phi_{\text{sol}}(x)) \langle \eta^2(x) \rangle \right] \\ &= M_{\text{cl}}(m_0) - \frac{m}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{\sqrt{k^2 + m^2}} = M_{\text{cl}}(m). \end{aligned} \quad (55)$$

The logarithmically divergent integral is exactly cancelled by the counterterm δM (36) after the renormalization of m .

The question we need to answer now is what is the value of $\langle \text{sol}|H|\text{sol} \rangle$? We recall that our calculation of the soliton mass was aimed at finding the dependence of M , or, $\langle \text{sol}|H|\text{sol} \rangle$ on m , the renormalized mass. We were not able to determine a constant term in the unrenormalized $\langle \text{sol}|H|\text{sol} \rangle$, but we knew that it must be subtracted to satisfy the renormalization condition $M|_{m=0} = 0$. To evaluate the l.h.s. of (51) we have to know this constant. In order to find it we must evaluate directly the sums of bosonic and fermionic frequencies in the soliton sector. Although the sums are quadratically divergent supersymmetry improves the situation. It requires that bosonic and fermionic modes come in pairs. Therefore we need to apply the Euler-Maclaurin formula to a function $f(n) = (\omega_n^B - \omega_n^F)/2$ which has much better behavior at large n . Using the spectral relations for the bosons (9) and the fermions (33) we find:

$$\begin{aligned} \langle \text{sol}|H|\text{sol} \rangle &= M_{\text{cl}}(m_0) + \frac{1}{2} \sum_{n=-N}^N (\omega_n^B - \omega_n^F) = M_{\text{cl}}(m_0) + \frac{1}{2} \int_{-\Lambda}^{\Lambda} \frac{dk}{2\pi} \frac{\theta(k)}{2} \frac{d}{dk} \sqrt{k^2 + m^2} \\ &= M_{\text{cl}}(m_0) + \frac{1}{8\pi} \sqrt{k^2 + m^2} \theta(k) \Big|_{-\Lambda}^{\Lambda} - \frac{1}{4} \int_{-\Lambda}^{\Lambda} \frac{dk}{2\pi} \sqrt{k^2 + m^2} \theta'(k) \\ &\stackrel{\Lambda \rightarrow \infty}{=} M_{\text{cl}}(m_0) + \frac{\Lambda}{4} - \frac{m}{2\pi} - \frac{1}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{m}{\sqrt{k^2 + m^2}}. \end{aligned} \quad (56)$$

The constant Λ is the ultraviolet cutoff related to the number of modes: $L\Lambda = 2\pi(N+1/2) + O(1/L)$, and we recall that $\theta = -2 \arctan(m/k)$. The last integral is logarithmically divergent and is exactly cancelled by the counter-term δM (36) when we renormalize $m_0 = m + \delta m$. The same divergence appears in $\langle \text{sol}|Z|\text{sol} \rangle$ and is also removed when m is renormalized. On the other hand, the linear divergent term $\Lambda/4$ does not appear in $\langle \text{sol}|Z|\text{sol} \rangle$. In terms of the renormalized mass m and $M_{\text{cl}}(m)$ the left- and the right-hand sides of (51) are:

$$\langle \text{sol}|H|\text{sol} \rangle = M_{\text{cl}} + \frac{\Lambda}{4} - \frac{m}{2\pi} \quad \text{and} \quad \langle \text{sol}|Z|\text{sol} \rangle = M_{\text{cl}}. \quad (57)$$

We see that the bound is observed by the soliton vacuum state with an infinite overkill due to a linearly divergent constant $\Lambda/4$. This constant is only nonzero in the soliton sector. In this sector the bound is not saturated, which shows that “what can happen — does happen”: the argument of Olive and Witten for the saturation of the bound is based on the “multiplet shortening” and does not apply to $N = 1$ susy solitons.

In other words, to resolve the long-standing problem of the Bogomolnyi bound in the $N = 1$ supersymmetric soliton/kink model we must realize that the bound is imposed on the unrenormalized expectation value of the Hamiltonian $\langle \text{sol} | H | \text{sol} \rangle$. Both sides of equation (51) contain ultraviolet divergences. The supersymmetry ensures that quadratic divergences do not appear in $\langle H \rangle$. However, $N = 1$ is not enough and a linear divergence remains in the topologically nontrivial sector (there is also a logarithmic divergence, but it is matched on both sides, $\langle H \rangle$ and $\langle Z \rangle$). This linearly divergent term is positive in accordance with the bound (51). Note that this divergence is different from a cosmological constant (which vanishes because of supersymmetry) in that it is proportional to L^0 rather than L^1 .

Since $\langle \text{sol} | H | \text{sol} \rangle$ is divergent even after standard renormalization of the mass m we need to use an additional renormalization condition to find the physical soliton mass. This is the condition $M|_{m=0} = 0$ which we introduced. Therefore

$$M = \langle \text{sol} | H | \text{sol} \rangle - \langle \text{sol} | H | \text{sol} \rangle|_{m=0} = \langle \text{sol} | H | \text{sol} \rangle - \frac{\Lambda}{4}. \quad (58)$$

Our new renormalization condition is based on the physical requirement that the physical vacuum energy should not depend on topology in the conformal point $m = 0$. Therefore, in principle, it also requires a subtraction of the expectation value of H in the topologically trivial vacuum $\langle 0 | H | 0 \rangle$, or rather $\langle 0 | H | 0 \rangle - \langle 0 | H | 0 \rangle|_{m=0}$. In the non-supersymmetric case this is essential to cancel the background bulk contributions linear in L , which are m -dependent, and which are the same in both sectors. However, supersymmetry ensures that $\langle 0 | H | 0 \rangle = 0$. Finally, we find a negative finite quantum correction to the physical soliton mass given by (37). However, it is not the physical mass M , but rather it is $\langle \text{sol} | H | \text{sol} \rangle$, to which the bound (51) applies. As we shall see in the next section, in the case of $N = 2$ supersymmetry all corrections, even finite, are cancelled and $M = \langle \text{sol} | H | \text{sol} \rangle$ saturates the bound as the “multiplet shortening” demands.

5 The N=2 case

Consider the action for the following N=(2,2) susy model in 1+1 dimensions

$$\mathcal{L} = -\partial_\mu \varphi^* \partial^\mu \varphi - \bar{\psi} \gamma^\mu \partial_\mu \psi - U^* U - \frac{1}{2} (U' \psi^T i \gamma^0 \psi + U^{*'} \psi^{*T} i \gamma^0 \psi^*) \quad (59)$$

where φ and ψ are complex, $U = U(\varphi)$, $U' = (\partial/\partial\varphi)U$, and $\gamma^1 = \tau_3$ and $\gamma^0 = -i\tau_2$ with τ_3 and τ_2 the usual Pauli matrices.⁵ The action is invariant under $\delta\varphi = \bar{\epsilon}\psi$, $\delta\varphi^* = \bar{\psi}\epsilon$, $\delta\psi = \not{\partial}\varphi\epsilon - U^*\epsilon^*$ and $\delta\bar{\psi} = -\bar{\epsilon}\not{\partial}\varphi^* - U\epsilon^T i\gamma^0$ with complex two-component spinors ϵ , with $\bar{\epsilon} = \epsilon^\dagger i\gamma^0$ and $\bar{\psi} = \psi^\dagger i\gamma^0$.

⁵With $\gamma^1 = \tau_3$ instead of $\gamma^1 = \tau_1$ the diagonalization of the fermionic actions is easier.

In terms of the components $\varphi = (\varphi_1 + i\varphi_2)/\sqrt{2}$ and $\psi^T = (\psi_+, \psi_-)$ the action reads

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\partial_\mu\varphi_1\partial^\mu\varphi_1 - \frac{1}{2}\partial_\mu\varphi_2\partial^\mu\varphi_2 + i\psi_+^*(\dot{\psi}_+ - \psi'_-) + i\psi_-^*(\dot{\psi}_- - \psi'_+) \\ & -U^*U + iU'\psi_+\psi_- + iU'^*\psi_+^*\psi_-^* \end{aligned} \quad (60)$$

For the superkink

$$U_{\text{kink}} = \sqrt{\lambda}(\varphi^2 - \mu_0^2/2\lambda) \quad (61)$$

while for super sine-Gordon theory

$$U_{\text{sine-Gordon}} = m_0^2\sqrt{2/\lambda}\cos(\sqrt{\lambda/2}\phi/m_0) \quad (62)$$

In terms of φ_1 and φ_2 the potential is given by

$$U^*U_{\text{kink}} = \frac{\lambda}{4} \left[\left(\varphi_1^2 - \mu_0^2/\lambda \right)^2 + 2\varphi_1^2\varphi_2^2 + \varphi_2^4 + 2\frac{\mu_0^2}{\lambda}\varphi_2^2 \right] \quad (63)$$

$$U^*U_{\text{sine-Gordon}} = \frac{m_0^4}{\lambda} \left[\cos \frac{\sqrt{\lambda}}{m_0}\varphi_1 + \cosh \frac{\sqrt{\lambda}}{m_0}\varphi_2 \right] \quad (64)$$

This already shows that for the kink the trivial solutions $\varphi_1^{(0)} = \pm\mu_0/\sqrt{\lambda}$ and the kink solution $\varphi_1 = \varphi_K = (\mu_0/\sqrt{\lambda})\tanh\mu_0 x/\sqrt{2}$ (and the antikink solution $\varphi_{\bar{K}} = -\varphi_K$) of the bosonic kink model remain solutions of this susy model while $\varphi_2 = 0$. Because the potential is of the form $(\varphi^2 - \mu_0^2/2\lambda)(\varphi^{*2} - \mu_0^2/2\lambda)$ instead of $(\varphi\varphi^* - \mu_0^2/2\lambda)^2$, there is no $U(1)$ symmetry acting on (φ_1, φ_2) which can rotate the kink away. Hence there is a genuine soliton.

In sine-Gordon theory, the trivial vacua are at $\varphi_2^{(0)} = 0$ and $\varphi_1^{(0)} = (n + \frac{1}{2})\pi m/\sqrt{\lambda}$, while for the vacuum in the topological sector we choose the solution

$$\varphi_{\text{sol}}^{\text{SG}}(x) = m/\sqrt{\lambda}(4\arctan(\exp mx) - \pi).$$

This solution is antisymmetric in x , in agreement with the Z_2 topological boundary conditions of section 2.⁶

The transformation rules in component form read

$$\begin{aligned} \delta\varphi &= -i\epsilon_+^*\psi_- + i\epsilon_-^*\psi_+ & \delta\psi_+ &= \varphi'\epsilon_+ - \dot{\varphi}\epsilon_- - U^*\epsilon_+^* \\ \delta\varphi^* &= -i\psi_+^*\epsilon_- + i\psi_-^*\epsilon_+ & \delta\psi_- &= -\varphi'\epsilon_- + \dot{\varphi}\epsilon_+ - U^*\epsilon_-^* \end{aligned} \quad (65)$$

and since for the soliton $\varphi'_{\text{sol}} + \sqrt{2}U(\varphi_{\text{sol}}/\sqrt{2}) = 0$, the solution $\varphi = (1/\sqrt{2})\varphi_{\text{sol}}$, $\psi_+ = \psi_- = 0$ is only preserved by half the susy transformations, namely those with $Im\ \epsilon_+$ and $Re\ \epsilon_-$.

In the sector without soliton we set $\varphi_1 = \varphi_1^{(0)} + \eta$ and find then the following linearized field equations

$$\begin{aligned} \square\eta - m_0^2\eta &= 0, \quad \square\varphi_2 - m_0^2\varphi_2 = 0 \\ \left. \begin{aligned} \dot{\psi}_+ - \psi'_- + m_0\psi_-^* &= 0 \\ \dot{\psi}_- - \psi'_+ - m_0\psi_+^* &= 0 \end{aligned} \right\} \begin{aligned} \ddot{\psi}_+ - \psi_+'' + m_0^2\psi_+ &= 0 \\ \ddot{\psi}_- - \psi_-'' + m_0^2\psi_- &= 0 \end{aligned} \end{aligned} \quad (66)$$

⁶This solution is obtained from the one in [14] by the substitution $\varphi = \varphi' + m\pi/\sqrt{\lambda}$. Actually, the sine-Gordon model has Z symmetry, and we could choose solutions which interpolate between two other minima of the potential by constant shifts.

where $m_0 = U'(\phi_1^{(0)}/\sqrt{2})$ and we used that $U(\phi_1^{(0)}/\sqrt{2}) = 0$. Hence all fields satisfy the same second-order field equation in the trivial sector, with a common mass m_0 (for the kink $m_0^2 = 2\mu_0^2$).

In the sector with a soliton, we set $\varphi_1 = \varphi_{\text{sol}}(x) + \eta$, and find then the linearized field equations

$$\begin{aligned} \square\eta - (U'^2 + UU'')\Big|_{\varphi_{\text{sol}}} \eta &= 0 \quad ; \quad \dot{\psi}_+ - \psi'_- + U'\Big|_{\varphi_{\text{sol}}} \psi_-^* = 0 \\ \square\varphi_2 - (U'^2 - UU'')\Big|_{\varphi_{\text{sol}}} \varphi_2 &= 0 \quad ; \quad \dot{\psi}_- - \psi'_+ - U'\Big|_{\varphi_{\text{sol}}} \psi_+^* = 0 \end{aligned} \quad (67)$$

Decomposing $\psi_+ = \text{Re } \psi_+ + i\text{Im } \psi_+$ and similarly for ψ_- , the fermionic field equations split into one pair of equations which couple $\text{Re } \psi_+$ to $\text{Re } \psi_-$, and another pair which couple $\text{Im } \psi_+$ to $\text{Im } \psi_-$. Iteration and the relation $\varphi'_{\text{sol}} = -\sqrt{2}U$ lead to

$$\begin{aligned} \text{Re } \psi_+'' - \text{Re } \ddot{\psi}_+ - (U'^2 + UU'')\Big|_{\varphi_{\text{sol}}} \text{Re } \psi_+ &= 0, \text{ idem for } \text{Im } \psi_- \\ \text{Re } \psi_-'' - \text{Re } \ddot{\psi}_- - (U'^2 - UU'')\Big|_{\varphi_{\text{sol}}} \text{Re } \psi_- &= 0, \text{ idem for } \text{Im } \psi_+ \end{aligned} \quad (68)$$

Hence, the real triplet $\eta, \text{Re } \psi_+$ and $\text{Im } \psi_-$ satisfies the same field equation as the real scalar η and the real upper component ψ_+ of the Majorana fermion in the N=(1,1) model, whereas the real triplet $\varphi_2, \text{Im } \psi_+$ and $\text{Re } \psi_-$ satisfies the same field equation as the real spinor ψ_- in the N=(1,1) model [14].

In principle one can directly determine the discrete spectrum of φ_2 by solving the Schrödinger equation with $V = (U')^2 - UU''$ [24], but susy already gives the answer. In the N=(1,1) model, the spinors $u_{\pm}$ in $\psi_{\pm}(x, t) = u_{\pm}(x) \exp -i\omega t$ satisfy the coupled equations $(\partial_x + \tilde{U}')u_+ + i\omega u_- = 0$ and $(\partial_x - \tilde{U}')u_- + i\omega u_+ = 0$, where $\tilde{U} = \sqrt{2}U(\varphi = \varphi_1/\sqrt{2})$. Any solution for u_+ and u_- yields also a solution for φ_2 (with $\varphi_2 \sim u_-$), and any solution for φ_2 leads also to a solution for u_+ and u_- (with $u_- \sim \varphi_2$ and $u_+ \sim (\partial_x - \tilde{U}')\varphi_2$). Hence, there are as many bound states for φ_2 as for η , namely $\varphi_{2,B} \sim (\partial_x + \tilde{U}')\eta_B$. (The zero mode $u_+ \sim \varphi'_{\text{sol}}$ does not lead to a corresponding solution for u_- and φ_2 since $(\partial_x + \tilde{U}')\varphi'_{\text{sol}} = 0$). The discrete spectrum of the fermions in the N=(2,2) model is then as follows: in the sector with $\text{Re } \psi_+$ and $\text{Re } \psi_-$ there are one discrete state with zero energy ($\text{Re } \psi_+ \sim \varphi'_{\text{sol}}$) and bound states with energy ω_B ($\text{Re } \psi_+ \sim \eta_B$). In the sector with $\text{Re } \psi_-$ and $\text{Im } \psi_-$ the same normalizable solutions are found. Thus, as expected, the massive bosonic spectrum of small oscillations around the soliton background is equal to the corresponding fermionic spectrum, and consists of massive quartets. There are also one bosonic (for translations) and two fermionic zero modes (zero energy solutions of the linearized field equations). The latter are proportional to the nonvanishing susy variations $\delta\psi \sim \varphi'_{\text{sol}}\epsilon$, which are due to the susy parameters $\text{Re } \epsilon_+$ and $\text{Im } \epsilon_-$.⁷ The zero modes do not form a susy multiplet (there are two fermionic and only one bosonic zero mode) but this poses no problem as quantization of collective coordinates tells us that the translational zero mode does not correspond to a physical particle. The fermionic zero modes are due to translations in superspace, namely when the susy generators $Q_{\pm}$ act on the superfield $\Phi(x, \theta) = \varphi_{\text{sol}}$.

⁷One can directly determine these fermionic zero modes from (67) by looking for time-independent normalizable solutions. One finds then $\text{Re } \psi_+ \sim \text{Im } \psi_- \sim \exp[-\int_0^x U'(\phi_{\text{sol}}(x')/\sqrt{2})dx']$. These functions are indeed proportional to φ'_{sol} .

The topological boundary conditions for the action (59) are

$$\begin{aligned}\phi(-L/2) &= (-1)^p \phi(L/2); & \phi'(-L/2) &= (-1)^p \phi'(L/2); \\ \psi(-L/2) &= (-1)^q (\gamma_3)^p \psi(L/2).\end{aligned}\quad (69)$$

For the continuous spectrum with $p = 1$ and $q = 0$ or $q = 1$ boundary conditions, the quantization conditions are

$$\begin{aligned}\eta : & \quad kL + \delta(k) = 2n\pi + \pi \\ \phi_2 : & \quad kL + \delta(k) + \theta(k) = 2n\pi + \pi \\ \text{Re } \psi_+, \text{Im } \psi_+, \text{Re } \psi_-, \text{Im } \psi_- : & \quad kL + \delta(k) + \frac{1}{2}\theta(k) = 2n\pi + q\pi\end{aligned}\quad (70)$$

The one-loop corrections to the soliton mass, differentiated w.r.t. m , are then given by

$$\begin{aligned}\partial_m M^{(1)} &= \partial_m \delta M^{(1)} \\ &+ \frac{1}{2m} \int_{-\infty}^{\infty} \sqrt{k^2 + m^2} \left(\delta'(k) + \{\delta'(k) + \theta'(k)\} - 2\{\delta'(k) + \frac{1}{2}\theta'(k)\} \right) \frac{dk}{2\pi}\end{aligned}\quad (71)$$

where the massive bound states do not contribute because they come in susy multiplets. The continuum states do not contribute either, because all phase shifts clearly cancel. The counterterm δM vanishes in the $N = 2$ model since the η and ϕ_2 tadpole give $\frac{-3}{2}$ and $\frac{-1}{2}$ times a fermionic tadpole, respectively. Note that all these cancelations are in fact independent of any particular regularization scheme since all integrands cancel.

To decide whether the Bogomolnyi bound is saturated we now turn to the central charges. The super Poincaré charges are obtained by the Noether method and read

$$\begin{aligned}Q_+ &= \int (U\psi_- + \psi_+^* \dot{\varphi} + \psi_-^* \varphi') dx \\ Q_- &= - \int (U\psi_+ - \psi_-^* \dot{\varphi} - \psi_+^* \varphi') dx\end{aligned}\quad (72)$$

With topological boundary conditions the combination $Q_+ - Q_-$ is conserved, but not $Q_+ + Q_-$,

$$\dot{Q}_+ + \dot{Q}_- = -2 \left[U(\psi_+ - \psi_-) + (\psi_+^* + \psi_-^*)(\dot{\varphi} + \varphi') \right]_{L/2} \quad (73)$$

There are no ordering ambiguities in these operators, and using equal-time canonical commutation relations one finds the following algebra for $A_{\pm} = Q_{\pm} \pm (Q_{\pm})^*$ and $B_{\pm} = Q_{\pm} \pm Q_{\pm}^*$

$$\begin{aligned}\{A_{\pm}, A_{\pm}\} &= \pm 2H - (Z + Z^*); & \{A_+, A_-\} &= -(Z - Z^*) \\ \{B_{\pm}, B_{\pm}\} &= \pm 2H + 2\text{Re } Z; & \{B_+, B_-\} &= Z - Z^* \\ \{A_{\pm}, B_{\pm}\} &= \pm 2P; & \{A_+, B_-\} &= \{A_-, B_+\} = 0\end{aligned}\quad (74)$$

The generators which are produced on the right-hand side are

$$\begin{aligned}H &= \int_{-\infty}^{\infty} \left(\dot{\varphi} \dot{\varphi}^* + \varphi' \varphi'^* + UU^* + i\psi_+^* \dot{\psi}_+ + i\psi_-^* \dot{\psi}_- \right) dx \\ P &= \int_{-\infty}^{+\infty} \left(\varphi' \dot{\varphi}^* + \dot{\varphi} \varphi'^* + i\psi_+^* \psi'_+ + i\psi_-^* \psi'_- \right) dx \\ Z &= 2 \int_{-\infty}^{\infty} U \varphi' dx\end{aligned}\quad (75)$$

Since an $N = 1$ massless multiplet in $D = (3, 1)$ (which is always without central charge) becomes a massive $N = 2$ multiplet in $D = (1, 1)$ whose $(\text{mass})^2$ is equal to the square of the central charges, while the $N=(1,1)$ susy algebra in $D = (3, 1)$ becomes an $N=(2,2)$ susy algebra in $D = (1, 1)$ with two central charges (the generators P_2 and P_3), it is clear why massive multiplets of $D = (1, 1)$ $N = (2, 2)$ models with maximal central charge are shortened.

Since A_+ and iA_- , and B_+ and iB_- are hermitian, the BPS bound is $H \geq |Re Z|$. For the soliton one has classically that Z is real and $H = Re Z$, i.e., the bound is saturated. At the quantum level, the 1-loop corrections to $Re Z$ are given by expanding $\varphi = \varphi_{\text{sol}}/\sqrt{2} + \chi$ and taking the vacuum expectation values in the soliton vacuum

$$\begin{aligned} Z &= Z_{cl} + Re \int \left\{ U'(\varphi_{\text{sol}}/\sqrt{2}) \langle \chi \chi' \rangle + \frac{1}{2} U''(\varphi_{\text{sol}}/\sqrt{2}) \langle \chi \chi \rangle \varphi'_{\text{sol}}/\sqrt{2} \right\} dx \\ &= Z_{cl} + Re \int \partial_x \left[\frac{1}{2} U'(\varphi_{\text{sol}}/\sqrt{2}) \langle \chi \chi \rangle \right] dx \\ &= Z_{cl} + \frac{m}{2} Re (\langle \chi(+\infty) \chi(+\infty) \rangle - \langle \chi(-\infty) \chi(-\infty) \rangle) \end{aligned} \quad (76)$$

Since asymptotically $\langle \chi \chi \rangle = 0$ (only $\langle \chi^* \chi \rangle$ is nonzero), there is no correction to the central charges. Because there is also no correction to the mass of the soliton, the BPS bound remains saturated.

6 Higher loops

In this section we repeat the two-loop calculation of the mass of the soliton in the sine-Gordon theory [20, 19], paying close attention this time to possible ambiguities. We begin with a review of the method of quantization of collective coordinates, focusing on possible ordering ambiguities. For early work on quantization of collective coordinates see [25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35]. For an introduction see [1].

To compute the higher loop corrections to the mass of a soliton, one may use standard quantum mechanical perturbation theory. One expands the renormalized Hamiltonian into a free part and an interaction part, $H = H^0 + H_{\text{int}}$, and the latter is expanded in terms of the dimensionless interaction parameter $\sqrt{\hbar c \lambda / m^2}$ as $H_{\text{int}} = \sum_{n=1}^{\infty} H_{\text{int}}^{(n)}$.⁸ This expansion is performed both in the soliton sector and in the trivial sector, and the corrections to the mass of the soliton can then be evaluated to any given order in $\hbar \lambda / m^2$ by subtracting the energy of the vacuum in the trivial sector from the energy of the vacuum in the soliton sector. For the two-loop corrections (themselves of order $\hbar^2 \lambda / m$ since the classical energy M_{cl} of the soliton is proportional to m^3 / λ) this means that we must evaluate

$$\begin{aligned} M^{(2)} &= \langle \text{sol} | H_{\text{int}, \text{sol}}^{(2)} | \text{sol} \rangle - \langle 0 | H_{\text{int}, \text{triv}}^{(2)} | 0 \rangle \\ &\quad + \sum_p' \frac{\langle \text{sol} | H_{\text{int}, \text{sol}}^{(1)} | p \rangle \langle p | H_{\text{int}, \text{sol}}^{(1)} | \text{sol} \rangle}{E_{\text{sol}} - E_p} \end{aligned}$$

⁸Terms with n quantum fields η contain in addition a dimensionless factor $(\omega L / c)^{-n/2}$. We set $c = 1$ but keep $\hbar$ when useful.

$$-\sum_p \frac{\langle 0 | H_{\text{int, triv}}^{(1)} | p \rangle \langle p | H_{\text{int, triv}}^{(1)} | 0 \rangle}{E_0 - E_p} \quad (77)$$

Here $|\text{sol}\rangle$ is the ground state in the soliton sector (the soliton vacuum) with classical energy $E_{\text{sol}} = M_{cl}$, $|0\rangle$ is the ground state in the trivial (nontopological) sector with vanishing classical energy, $|p\rangle$ are the complete sets of eigenstates of $H_{\text{sol}}^{(0)}$ and $H_{\text{triv}}^{(0)}$ with positive energies E_p , and the sums extend over all excitations but do not include the ground state. Hence $E_{\text{sol}} - E_p$ and $E_0 - E_p$ never vanish. The fields for the quantum fluctuations in the trivial and the topological sectors are expanded into modes with creation and annihilation operators, and both $|0\rangle$ and $|\text{sol}\rangle$ are annihilated by the annihilation operators.

6.1 The quantum Hamiltonian

To apply this approach to the soliton in sine-Gordon theory, we begin by defining the sine-Gordon action⁹

$$\mathcal{L} = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}(\phi')^2 - \frac{m_0^4}{\lambda} \left(1 - \cos \frac{\sqrt{\lambda}}{m_0} \phi \right) \quad (78)$$

The action in the trivial sector is obtained by expanding ϕ about the trivial vacuum. Since for sine-Gordon theory the latter is given by $\Phi=0$, we obtain $\phi = \Phi + \eta = \eta$, and

$$\mathcal{L} = \frac{1}{2}\dot{\eta}^2 - \frac{1}{2}(\eta')^2 - \frac{1}{2}m_0^2\eta^2 + \frac{1}{4!}\lambda\eta^4 + \dots \quad (79)$$

In 1+1 dimensional linear sigma models only mass renormalization is needed, $m_0^2 = m^2 + \delta m^2$, and at the one-loop level δm^2 is fixed by requiring that the graph with a seagull loop and two external η fields cancels the contribution from $-(1/2)\delta m^2\eta^2$. This yields

$$\delta m^2 = \frac{\lambda}{2} \sum \frac{\hbar}{2\omega_{\text{vac}}L} = \frac{\hbar\lambda}{4\pi} \int_0^\Lambda \frac{dk}{\sqrt{k^2 + m^2}} \quad (80)$$

The mass m is thus the physical mass of the meson at the pole of the propagator to this order.

A complete counterterm which removes all equal-time contractions is [15]

$$\Delta H = (e^{\delta m^2/m^2} - 1) \frac{m^4}{\lambda} \int_{-\infty}^{+\infty} \left(1 - \cos \frac{\sqrt{\lambda}}{m} \phi(x) \right) dx \quad (81)$$

The corrections to the physical mass at higher loop orders are then finite, and by expanding the final result for the soliton mass in terms of the physical mass of the mesons, any ambiguity due to defining a ΔH which differs from (81) by finite terms will be eliminated. In particular, the contributions from other renormalization conditions for m , and finite renormalization of λ and η , should cancel.

⁹To compare with [19, 20] we use their action. It is related to the action in section 5 by the shift $\sqrt{\lambda}\varphi_1/m_0 \rightarrow \sqrt{\lambda}\varphi_1/m_0 + \pi$. Note that in [20] mass is renormalized by $m_0^2 = m^2 - \delta m^2$.

The Hamiltonian in the trivial sector is simply

$$H_{\text{triv}}^{(0)} = \int_{-\infty}^{+\infty} \left[\frac{1}{2} \Pi_0^2(x) + \frac{1}{2} (\eta'(x))^2 + \frac{1}{2} m^2 \eta(x)^2 \right] dx \quad (82)$$

$$H_{\text{int, triv}} = -\frac{1}{4!} \lambda \eta^4 + \dots + \frac{1}{2} \delta m^2 \eta^2 + \dots \quad (83)$$

where $\Pi_0(x)$ is the momentum canonically conjugate to $\eta(x)$. So $H_{\text{int, triv}}^{(n)}$ contains only terms for even n and to obtain the contributions of the trivial sector to the two-loop corrections to the mass of the soliton we must evaluate $-\frac{1}{4!} \lambda \langle 0 | \eta^4 | 0 \rangle + \frac{1}{2} \delta m^2 \langle 0 | \eta^2 | 0 \rangle$. Note that there are no ordering ambiguities in the Hamiltonian of the trivial sector.

To obtain the Hamiltonian in the soliton sector, one must use the formalism for quantization of collective coordinates [1]. Although the final formulas look somewhat complicated, the basic idea is very simple: one expands $\phi(x, t)$ again into a sum of a background field (the soliton) and a complete set of small fluctuations about the background field, but instead of simply writing $\phi(x, t) = \phi_{\text{sol}}(x) + \sum q^m(t) \eta_m(x)$ where $\eta_m(x)$ stands for all modes (eigenfunctions of the linearized field equations), one deletes the zero mode for translations from the sum, and reintroduces it by replacing x by $x - X(t)$ on the right hand side of the expansion of ϕ . For small $X(t)$, the expansion of $\phi_{\text{sol}}(x - X(t))$ into a Taylor series gives $\phi_{\text{sol}} - X(t) \phi'_{\text{sol}}(x, t) + \dots$, and since $\phi'_{\text{sol}}(x, t)$ is the translational zero mode (the solution of the linearized field equations with vanishing energy), one has not lost any degrees of freedom. Hence one substitutes

$$\phi(x, t) = \phi_{\text{sol}}(x - X(t)) + \sum' q^m(t) \eta_m(x - X(t)) \quad (84)$$

into the action in (78), and using the chain rule, one finds an action of the form of a quantum mechanical nonlinear sigma model (but with infinitely many degrees of freedom)

$$L = \dot{u}^I g_{IJ}(u) \dot{u}^J - V(u); u^I = \{X(t), q^m(t)\} \quad (85)$$

The metric g_{IJ} is given by

$$g_{IJ} = \int \frac{\partial \phi(x, t)}{\partial u^I} \frac{\partial \phi(x, t)}{\partial u^J} dx \quad (86)$$

and contains space integrals over expressions which depend on $q^m(t)$, $\eta_m(x)$ and $\phi_{\text{sol}}(x)$, but not on $X(t)$ due to the translational invariance of the integral over x . The Hamiltonian is then simply given by

$$H = \pi_I g^{IJ}(u) \pi_J + V(u); \pi_I = \{P(t), \pi_m(t)\} \quad (87)$$

where $g^{IJ}(u)$ is the matrix inverse of the metric $g_{IJ}(u)$ and $P(t)$ is the center of mass momentum (the momentum conjugate to $X(t)$), while $\pi_m(t)$ are momenta canonically conjugate to $q^m(t)$.

Classically, this is the whole result. One may check that the equal-time Poisson brackets $\{Q, P\} = 1$, $\{q^m, \pi_n\} = \delta_n^m$ imply $\{\phi(x), \Pi_0(y)\} = \delta(x - y)$ where $\Pi_0(x, t) = \dot{\phi}(x, t)$, and vice-versa. Hence, the transition from $\phi(x, t)$ and $\Pi_0(x, t)$ to $\{X(t), q^m(t)\}$ and $\{P(t), \pi_m(t)\}$ is a canonical transformation. It is useful to recast the “quantum mechanical” Hamiltonian

in (87) into a form which resembles more the Hamiltonian of a 1+1 dimensional field theory. To this purpose we introduce fields constructed from q^m and π_m as follows

$$\eta(x, t) \equiv \sum' q^m(t) \eta_m(x - X(t)) \quad (88)$$

$$\pi(x, t) \equiv \sum' \pi_m(t) \eta_m(x - X(t)) \quad (89)$$

By combining the π_m and q^m with the functions $\eta_m(x)$ which appear in $g^{IJ}(u)$, one can write the complete Hamiltonian only in terms of the fields $\eta(x, t)$ and $\pi(x, t)$ and the background field $\phi_{\text{sol}}(x)$. To simplify the notation, we introduce an inner product $(f, h) \equiv \int_{-\infty}^{+\infty} f^*(x) h(x) dx$. Note that the functions η_m which parameterize the small fluctuations are orthogonal to the zero mode ϕ'_{sol} since they correspond to different eigenvalues of the kinetic operator

$$(\phi'_{\text{sol}}, \eta_m) = 0 \quad (90)$$

a result we shall use repeatedly. The classical Hamiltonian density $\mathcal{H} = T_{00}$ is given by

$$\begin{aligned} \mathcal{H} &= \frac{1}{2} \Pi_0^2(x, t) + \frac{1}{2} \phi'(x, t)^2 + V(\phi) \\ &= \frac{1}{2} \pi^2(x - X(t), t) - \pi(x - X(t), t) \frac{P + (\pi, \eta')}{M_{cl} [1 + (\eta', \phi'_{\text{sol}})/M_{cl}]} \phi'_{\text{sol}}(x - X(t)) \\ &\quad + \frac{[P + (\pi, \eta')]^2}{2M_{cl}^2 (1 + (\eta', \phi'_{\text{sol}})/M_{cl})} [\phi'_{\text{sol}}(x - X(t))]^2 + \\ &\quad + \frac{1}{2} [\eta'(x - X(t), t) + \phi'_{\text{sol}}(x - X(t))]^2 + V(\phi_{\text{sol}} + \eta) \end{aligned} \quad (91)$$

A great simplification occurs in the Hamiltonian $H = \int_{-\infty}^{+\infty} \mathcal{H} dx$ because due to the orthogonality of the zero mode ϕ'_{sol} to the fluctuations η_m , the complicated second term in $\mathcal{H}$ cancels. There should be no terms linear in the fluctuations η and π and the collective coordinates X, P in H (i.e., after integrating $\mathcal{H}$ over x). That this is indeed the case follows from the field equation

$$\phi''_{\text{sol}} = V'(\phi_{\text{sol}}) \quad (92)$$

The classical energy of the soliton at rest is given by

$$M_{cl} = \int_{-\infty}^{+\infty} (\phi'_{\text{sol}})^2 dx \quad (93)$$

which follows from equipartition of energy

$$\frac{1}{2} (\phi'_{\text{sol}})^2 = V(\phi_{\text{sol}}) \quad (94)$$

Thus we arrive at the following expression for the classical Hamiltonian in the topological sector

$$H_{\text{sol}}^{(0)} = M_{cl} + \int_{-\infty}^{+\infty} \left[\frac{1}{2} \pi(x, t)^2 + \frac{1}{2} \eta'(x, t)^2 + \frac{1}{2} \eta^2 V''(\phi_{\text{sol}}) \right] dx \quad (95)$$

$$\begin{aligned} H_{\text{int, sol}}^{\text{cl}} &= \frac{1}{2M_{cl}} \frac{[P + (\pi, \eta')]^2}{1 + (\eta', \phi'_{\text{sol}})/M_{cl}} + \int \left[\frac{1}{3!} \eta^3 V'''(\phi_{\text{sol}}) + \right. \\ &\quad \left. + \frac{1}{4!} \eta^4 V''''(\phi_{\text{sol}}) + \dots \right] dx \end{aligned} \quad (96)$$

All $X(t)$ dependence has disappeared from H due to translational invariance of the integration over x .

We must now discuss the subtle issue of operator ordering in H . We shall consider a soliton at rest, so we set $P = 0$. Furthermore, due to $[q^m, \pi_n] = i\hbar\delta_n^m$ and $(\eta_m, \eta'_m) = 0$ (since we work in a finite volume, and η and η' have the same boundary conditions) one has the equality $(\pi, \eta') = (\eta', \pi)$, at least if one considers a finite number of modes in η and π . However, there are operator ordering ambiguities both in $(\pi, \eta')^2$ and also with respect to the term $(\eta', \phi'_{\text{sol}})/M_{\text{cl}}$ in the denominator.

In general, one may require that the generators $H, P = \int T_{01}dx$ and $L = \int xT_{00}dx$ satisfy the Poincaré algebra [33]¹⁰. The expressions for these operators are quite complex, and in general it seems likely that the operator ordering which leads to closure of the Poincaré algebra is unique (in quantum gravity, such an ordering has never been found). There is, however, an ordering which guarantees closure, and this is the ordering we shall adopt. It is obtained by making the canonical transformation at the quantum level. One begins with the quantum Hamiltonian in “Cartesian coordinates” (i.e., in terms of the operators $\Pi_0(x)$ and $\phi(x)$). In the Schrödinger representation the operator $\Pi_0(x)$ is represented by $\partial/\partial\phi(x)$, and making the change of coordinates from $\phi(x)$ to X and q^m , one obtains the Laplacian in curved space by applying the chain rule

$$\sum \left(\frac{\partial}{\partial \alpha^i} \right)^2 = \frac{1}{\sqrt{g}} \frac{\partial}{\partial u^I} \sqrt{g(u)} g^{IJ}(u) \frac{\partial}{\partial u^J} \quad (97)$$

where α^i is the set $\phi(x)$ and u^I the set X, q^m . If the inner product in α space is given by $(f, h) = \int f^*(\alpha) h(\alpha) (\prod d\alpha^i)$, it becomes in u space $(f, h) = \int f^*(\alpha(u)) h(\alpha(u)) \sqrt{g(u)} (\prod du^I)$. With this inner product, the relation between $\partial/\partial u^J$ and the conjugate momenta π_J in the Schrödinger representation is not simply $\pi_J = \partial/\partial u^J$, but rather

$$\frac{\partial}{\partial u^J} = g^{1/4}(u) \pi_J g^{-1/4}(u) \quad (98)$$

as one may check.¹¹ Hence the correct quantum Hamiltonian is given by

$$\hat{H} = \frac{1}{2} \frac{1}{g(u)^{1/4}} \pi_I \sqrt{g(u)} g^{IJ}(u) \pi_J \frac{1}{g(u)^{1/4}} + V(u) + \Delta H \quad (99)$$

with ΔH given by (81).

It is often useful to *rewrite* this Hamiltonian such that all expressions are Weyl ordered, because then one can use Berezin’s theorem and find at once the action to be used in the

¹⁰The Noether current for the orbital part of the angular momentum $J_{\rho\sigma}$ is given by $j_{\rho\sigma}^\mu = (x_\rho T_\sigma^\mu - x_\sigma T_\rho^\mu)$ so $J_{01} = L = \int (x_0 T_1^0 - x_1 T_0^0) dx$. At $t=0$ this reduces to $\int x T_{00} dx$.

¹¹In [1] this derivation of the operator ordering of the Hamiltonian is given, but at the end $\partial/\partial u^J$ is replaced by π_J which is incorrect. In [34] the correct quantum Hamiltonian is obtained, but the factors $g^{-1/4}(u)$ are produced by “Redefining the Hilbert space so as to eliminate the measure from this scalar product...”. We claim that the relation (98) is not a convention or a choice of basis, but is fixed because the inner product has been specified.

path integral. The result is

$$\hat{H} = \frac{1}{2}(\pi_I g^{IJ} \pi_J)_W + V(u) + \Delta V + \Delta H \quad (100)$$

$$\Delta V = \frac{\hbar^2}{8} \left[\partial_I \partial_J g^{IJ}(u) - 4g^{-1/4}(u) \partial_I \{g^{1/2}(u) g^{IJ}(u) \partial_J g^{-1/4}(u)\} \right] \quad (101)$$

The operator $(1/2)(\pi_I g^{IJ} \pi_J)_W$ is obtained by promoting (95) and (96) to operators and Weyl ordering. Weyl ordering yields then $(1/2)((1/4)\pi_I \pi_J g^{IJ} + (1/2)\pi_I g^{IJ} \pi_J + (1/4)g^{IJ} \pi_I \pi_J)$. Substituting the expression for $g^{IJ}(u)$ [1] one finds

$$\begin{aligned} \partial_I \partial_J g^{IJ} &= \partial_{q^m} \partial_{q^n} \left\{ \frac{(\eta^m, \eta')(\eta^n, \eta')}{(\psi_0, \phi'_{\text{sol}} + \eta')^2} \right\} \\ 4g^{-1/4} \partial_I \{g^{1/2} g^{IJ} \partial_J g^{-1/4}\} &= \frac{1}{(\psi_0, \phi')^{1/2}} \frac{\partial}{\partial q^m} \left[\frac{1}{(\psi_0, \phi')^{3/2}} \frac{\partial}{\partial q^m} (\psi_0, \phi')^2 + \right. \\ &\quad \left. + \frac{(\eta', \eta_m)(\eta', \eta_n)}{(\psi_0, \phi')^{7/2}} \frac{\partial}{\partial q^n} (\psi_0, \phi')^2 \right] \end{aligned} \quad (102)$$

where $\phi = \phi_{\text{sol}} + \eta$, and $\psi_0 = \phi'_{\text{sol}}/\sqrt{M_{\text{cl}}}$ is the normalized zero mode. This leads to

$$\begin{aligned} \Delta V &= \frac{\hbar^2}{8} \left[-\frac{(\psi_0, \eta'_m)(\eta'_m, \psi_0)}{(\psi_0, \phi')^2} \right. \\ &\quad - 2 \frac{(\psi_0, \eta'_m)(\eta_m, \eta'_n)(\eta_n, \eta') + (\psi_0, \eta'_m)(\eta_m, \eta')(\eta_n, \eta'_n)}{(\psi_0, \phi')^3} \\ &\quad \left. + \frac{\{(\psi_0, \eta'_m)(\eta_m, \eta')\}^2}{(\psi_0, \phi')^4} + \frac{(\eta_m, \eta'_n)(\eta_n, \eta'_m) + (\eta_m, \eta'_m)^2}{(\psi_0, \phi')^2} \right] \end{aligned} \quad (103)$$

Further simplifications result by using the identities

$$\begin{aligned} (\psi_0, \psi'_0) &= 0, \quad (\psi_0, \eta') = (\psi_0, \phi'), \quad (\psi_0, \eta'_m) = -(\psi'_0, \eta_m) \\ (\eta_n, \eta') &= (\eta_n, \phi'), \quad (\eta_m, \eta'_m) = 0, \quad \sum' \eta_m(x) \eta_m(y) = \delta(x - y) - \psi_0(x) \psi_0(y) \end{aligned} \quad (104)$$

The final answer for ΔV reads then

$$\begin{aligned} \Delta V &= \frac{\hbar^2}{8} \left[-\frac{(\psi'_0, \psi'_0)}{(\psi_0, \phi')^2} + 2 \left\{ \frac{(\psi'_0, \phi'')}{(\psi_0, \phi')^3} - \frac{(\psi'_0, \psi'_0)}{(\psi_0, \phi')^2} \right\} \right. \\ &\quad \left. + \frac{(\psi'_0, \phi')^2}{(\psi_0, \phi')^4} - \sum_{m,n} \frac{|(\eta_m, \eta'_n)|^2}{(\psi'_0, \phi')^2} \right] \end{aligned} \quad (105)$$

The total Hamiltonian is then the sum of $H_{\text{sol}}^{(0)}$ in (95) in which no ordering problems are present, and $(H_{\text{int,sol}}^{\text{cl}})_W + \Delta V + \Delta H$ with $(H_{\text{int,sol}}^{\text{cl}})_W$ given by (96) with the complicated momentum dependent term Weyl-ordered. This is the result in [34]. A drastic simplification is obtained by rewriting the latter term in a particular non-Weyl-ordered way in such a way

that it absorbs all terms in ΔV except the first one [33]. This leads to the final form of the interaction Hamiltonian¹²

$$\begin{aligned}
H_{\text{int,sol}} = & \frac{1}{8M_{\text{cl}}} \left\{ (P + (\eta', \pi)), \frac{1}{1 + (\eta', \phi'_{\text{sol}})/M_{\text{cl}}} \right\}^2 \\
& - \frac{\hbar^2}{8M_{\text{cl}}^2} \int_{-\infty}^{+\infty} \frac{(\phi''_{\text{sol}})^2 dx}{[1 + (\eta', \phi'_{\text{sol}})/M_{\text{cl}}]^2} + \Delta H \\
& + \int_{-\infty}^{+\infty} \left[\frac{1}{3!} \eta^3 V'''(\phi_{\text{sol}}) + \frac{1}{4!} \eta^4 V''''(\phi_{\text{sol}}) + \dots \right] dx
\end{aligned} \tag{106}$$

Note that the first term is the square of a Weyl-ordered operator, but is not itself Weyl-ordered.

For the two-loop calculation we are going to perform we set $P=0$, and we only need the terms as far as quartic in η and π . This leads to

$$\begin{aligned}
H_{\text{int,sol}} = & \frac{1}{2M_{\text{cl}}} \left(\int_{-\infty}^{+\infty} \eta' \pi dx \right) \left(\int_{-\infty}^{+\infty} \eta' \pi dy \right) - \frac{\hbar^2}{8M_{\text{cl}}^2} \int_{-\infty}^{+\infty} (\phi''_{\text{sol}}(x))^2 dx \\
& + \Delta H - \frac{\sqrt{\lambda} m}{3!} \int_{-\infty}^{+\infty} \eta^3(x) \sin \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) dx \\
& - \frac{\lambda}{4!} \int_{-\infty}^{+\infty} \eta^4(x) \cos \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) dx + \dots
\end{aligned} \tag{107}$$

The counterterms are the same in the topological sector as in the trivial sector, but in the trivial sector we decomposed $\phi = \Phi + \eta = \eta$, while here we expand $\phi = \phi_{\text{sol}} + \eta$. We obtain then

$$\begin{aligned}
\Delta H = & \frac{m^4}{\lambda} \left\{ e^{\frac{\delta m^2}{m^2}} - 1 \right\} \int_{-\infty}^{+\infty} \left\{ 1 - \cos \frac{\sqrt{\lambda}}{m} \phi(x) \right\} dx \\
= & \frac{m^4}{\lambda} \left[\frac{\delta m^2}{m^2} + \frac{1}{2} \left(\frac{\delta m^2}{m^2} \right)^2 \right] \int_{-\infty}^{+\infty} \left\{ 1 - \cos \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) \right\} dx \\
& + \frac{\delta m^2 m}{\sqrt{\lambda}} \int_{-\infty}^{+\infty} \eta(x) \sin \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) dx \\
& + \frac{1}{2} \delta m^2 \int_{-\infty}^{+\infty} \eta^2(x) \cos \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) dx + \dots
\end{aligned} \tag{108}$$

¹² The first term in (106) can be written for $P = 0$ as $[(\pi, \eta'/F) + (\eta'/F, \pi)]^2$, where $F = 1 + (\eta', \phi'_{\text{sol}})/M_{\text{cl}}$. Weyl-ordering of $(\pi, \eta'/F)^2$ yields

$$\frac{1}{4} \int_{-\infty}^{+\infty} \left[\pi(x) (\pi, \eta'/F) \frac{\eta'}{F}(x) + \frac{\eta'}{F}(x) (\eta'/F, \pi) \pi(x) \right] dx + 2(\pi, \eta'/F)(\eta'/F, \pi)$$

Evaluating the difference of the two expressions one needs the following commutators

$$\begin{aligned}
[\pi(x), \eta'(y)] &= -i \partial_y \delta(x - y) + \frac{i}{M_{\text{cl}}} \phi'_{\text{sol}}(x) \phi''_{\text{sol}}(y) \\
[\pi(x), 1/F] &= -i \phi''_{\text{sol}}(x) / (M_{\text{cl}} F^2)
\end{aligned}$$

Straightforward algebra produces then all terms in ΔV except the first one.

The first term is the counterterm for the one-loop graphs and will not contribute to our two-loop calculation.

6.2 The actual two-loop calculation

Using the explicit expressions for the classical soliton solution

$$\phi_{\text{sol}} = \frac{4m}{\sqrt{\lambda}} \arctan(e^{mx}) \quad (109)$$

one finds $M_{\text{cl}} = \frac{8m^3}{\lambda}$ and

$$\begin{aligned} \sin\left(\frac{\sqrt{\lambda}}{m}\phi_{\text{sol}}\right) &= -4\frac{e^{mx} - e^{-mx}}{(e^{mx} + e^{-mx})^2}; \quad \cos\frac{\sqrt{\lambda}}{m}\phi_{\text{sol}} = 1 - \frac{8}{(e^{mx} + e^{-mx})^2} \\ &\quad - \frac{1}{8M_{\text{cl}}^2} \int_{-\infty}^{+\infty} (\phi_{\text{sol}}''(x))^2 dx = -\frac{\lambda}{192m} \end{aligned} \quad (110)$$

Substituting these results into (107,108), we find for the Hamiltonian

$$H_{\text{int,sol}}^{(1),I} = -\frac{\sqrt{\lambda}m}{6} \int_{-\infty}^{+\infty} \frac{(-4)(e^{mx} - e^{-mx})}{(e^{mx} + e^{-mx})^2} \eta^3(x) dx \quad (111)$$

$$H_{\text{int,sol}}^{(1),II} = \frac{\delta m^2 m}{\sqrt{\lambda}} \int_{-\infty}^{+\infty} \frac{(-4)(e^{mx} - e^{-mx})}{(e^{mx} + e^{-mx})^2} \eta(x) dx \quad (112)$$

$$H_{\text{int,sol}}^{(2),I} = -\frac{\lambda}{24} \int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2}\right] \eta^4(x) dx \quad (113)$$

$$H_{\text{int,sol}}^{(2),II} = \frac{1}{2} \delta m^2 \int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2}\right] \eta^2(x) dx \quad (114)$$

$$H_{\text{int,sol}}^{(2),III} = \frac{\lambda}{16m^3} \left(\int_{-\infty}^{+\infty} \eta'(x) \pi(x) dx \right)^2 \quad (115)$$

$$H_{\text{int,sol}}^{(2),\text{rest}} = -\frac{\lambda \hbar^2}{192m} + \frac{(\delta m^2)^2}{2\lambda} \frac{4}{m} \quad (116)$$

We now put the system in a box of length L . We expand $\eta(x)$ into creation and annihilation operators

$$\eta(x) = \sum_{n=-\infty}^{+\infty} \sqrt{\frac{\hbar}{2\omega_n}} (a(q_n) E(q_n, x) e^{ik_n x} + h.c.) \quad (117)$$

$$E(q_n, x) = \frac{i \tanh mx + q_n}{[(1 + q_n^2)L - \frac{2}{m} \tanh \frac{mL}{2}]^{1/2}} \quad (118)$$

$$q_n = \frac{k_n}{m}; \quad \omega_n^2 = k_n^2 + m^2 \quad (119)$$

The functions $E(q_n, x) e^{ik_n x - i\omega_n t} \equiv \eta_n(x) e^{-i\omega_n t}$ are eigenfunctions of $H_{\text{sol}}^{(0)}$ and satisfy the linearized field equations

$$[\partial_x^2 + \omega_n^2 - m^2(1 - 2 \cosh^{-2}(x))] \eta_n = 0 \quad (120)$$

and for large q_n they tend to $\frac{1}{\sqrt{L}}$ which leads to the familiar normalization factor $(2\omega_n L)^{-1/2}$ for free fields. At $x = \pm L/2$ we find the phase shift

$$\delta(k) = 2 \arctan\left\{\frac{m}{k} \tanh \frac{1}{2}mL\right\} \quad (121)$$

The momenta k_n are discretized by adopting antiperiodic boundary conditions¹³

$$k_n L + \delta(k_n) = 2\pi n + \pi \quad (122)$$

As eigenfunctions of a self-adjoint operator on a compact space, the functions $E(q_n, x)$ have the same orthogonality properties as plane waves and they are normalized to unity, for example $\int_{-L/2}^{+L/2} |E(q_n, x)|^2 dx = 1$. As in the case of the kink $\delta(k)$ in (121) is discontinuous. The values of $n = \dots, -2, -1, 0, 1, 2, \dots$ give solutions of (122) $kL = \dots, -2\pi, 0, 0, 2\pi, 4\pi, \dots$. Again, we see a defect in the n to k mapping: the $k = 0$ solution is obtained for $n = -1$ and $n = 0$. If we map $k = 0$ onto $n = -1$, the remaining number $n = 0$ can be assigned to the only discrete solution (the translational zero mode). Then using the Euler-Maclaurin formula the sum over n can be converted into integral over k with continuous measure. We now evaluate the various terms in (77), adding subsets of terms which combine to cancel divergences.

The η^4 , $\delta m^2 \eta^2$, and $(\delta m^2)^2$ terms in $H_{\text{int,sol}}^{(2)}$ and $H_{\text{int,vac}}^{(2)}$

Since there are 3 ways to contract the 4 η 's, one finds from the term with η^4

$$\begin{aligned} \langle \text{sol} | H_{\text{int,sol}}^{(2),I} | \text{sol} \rangle &= -\frac{\lambda}{8} \int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2} \right] \sum_{n,p=-\infty}^{+\infty} \\ &\frac{(\tanh^2 mx - 1) + (q_n^2 + 1)}{(1 + q_n^2)L - \frac{2}{m} \tanh \frac{mL}{2}} \frac{(\tanh^2 mx - 1) + (q_p^2 + 1)}{(1 + q_p^2)L - \frac{2}{m} \tanh \frac{mL}{2}} \frac{dx}{2\omega_n 2\omega_p} \end{aligned} \quad (123)$$

Using the integrals in the appendix, we record the contributions due to terms with none, one and two factors $\tanh^2 x - 1$ in 3 separate lines

$$\begin{aligned} &= -\frac{\lambda}{32} \left[\frac{1}{L} \left(\sum \frac{1}{\omega} \right)^2 - \frac{4}{mL^2} \left(\sum \frac{1}{\omega} \right)^2 + \left\{ \frac{4}{L^2} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\omega^2} \right) + \dots \right\} \right. \\ &\quad + \frac{4}{3} \frac{m}{L^2} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\omega^3} \right) + \left\{ \frac{8}{3} \frac{1}{L^3} \left(\sum \frac{1}{\omega^3} \right)^2 + \frac{8}{3} \frac{1}{L^3} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\omega^5} \right) + \dots \right\} \\ &\quad \left. - \frac{4m^3}{5L^2} \left(\sum \frac{1}{\omega^3} \right)^2 + \left\{ -\frac{16m^2}{5L^3} \left(\sum \frac{1}{\omega^3} \right) \left(\sum \frac{1}{\omega^5} \right) + \dots \right\} \right] \end{aligned} \quad (124)$$

The first two terms come from the factor $1 - 8/(e^{mx} + e^{-mx})^2$ whose x -integral is equal to $L - \frac{4}{m}$. Clearly, there is a divergence proportional to L which will be cancelled by the corresponding contribution from the vacuum sector. The integrals of $1 - 8/(e^{mx} + e^{-mx})^2$ times $(\tanh^2 x - 1)^p$ for $p = 1$ and $p = 2$ are given by $\frac{2}{3m}$ and $-\frac{4}{5m}$, respectively, and do

¹³Of course, we differ here from refs. [19, 20] who choose periodic boundary conditions. The necessity for the antiperiodic boundary conditions has been discussed at length in the preceding sections.

not contain divergences which are due to the x integral, but they still contain divergences due to the sums over n and p . The terms inside curly brackets are due to expanding the denominators $(1 + q_n^2)L - \frac{2}{m} \tanh \frac{mL}{2} \sim (1 + q_n^2)L - \frac{2}{m}$. (We have already set $\tanh \frac{mL}{2} = 1$ since the difference vanishes exponentially fast for $L \rightarrow \infty$). Not all these terms vanish for $L \rightarrow \infty$, but they will cancel with similar terms from other contributions. The sums

$$\sum \frac{1}{\omega} \equiv \sum_{-\infty}^{+\infty} \frac{1}{\omega_n} \sim \frac{L}{2\pi} 2 \int_0^\infty \frac{1}{\omega(k)} \left[1 + \frac{1}{L} \delta'(k) \right] dk \quad (125)$$

will get contributions from $\delta'(k)$, but by first combining such sums, we shall find that only differences like $\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}}$ occur, and this will simplify the analysis significantly. We wrote an approximation symbol $\sim$ in $(\sum \frac{1}{\omega})$ because the evaluation of such sums has been found to be regularization dependent at one-loop level. This is the issue we want to study now at the two-loop level.

The evaluation of $\langle \text{sol} | H_{\text{int,sol}}^{(2),II} | \text{sol} \rangle$ is straightforward and yields

$$\begin{aligned} & \frac{1}{2} \delta m^2 \int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2} \right] \sum_n \frac{(\tanh^2 mx - 1) + (q_n^2 + 1)}{(q_n^2 + 1)L - \frac{2}{m}} \frac{dx}{2\omega_n} \\ &= \frac{1}{2} \delta m^2 \left(L - \frac{4}{m} \right) \left(\frac{1}{2L} \sum \frac{1}{\omega} + \left\{ \frac{1}{mL^2} \sum \frac{1}{\omega^3} + \dots \right\} \right) \\ & \quad + \frac{1}{2} \delta m^2 \left(\frac{2}{3m} \right) \frac{m^2}{2} \frac{1}{L} \sum \frac{1}{\omega^3} \end{aligned} \quad (126)$$

where the last line contains the contributions from the terms with $\tanh^2 x - 1$.

The contributions from the trivial sector are only $\frac{1}{2} \delta m^2 \langle 0 | \eta^2 | 0 \rangle - \frac{\lambda}{4!} \langle 0 | \eta^4 | 0 \rangle$, where $\eta(x)$ are now simple plane waves and they yield

$$\langle 0 | H_{\text{triv}} + \Delta H_{\text{triv}} | 0 \rangle = -\frac{1}{2} \delta m^2 \sum \frac{1}{2\tilde{\omega}} - \frac{\lambda}{8} \left(\sum \frac{1}{2\tilde{\omega}} \right)^2 \frac{1}{L} \quad (127)$$

Finally there is the contribution from the term proportional to $(\delta m^2)^2$ in ΔH_{sol} ; it yields

$$\langle \text{sol} | \frac{1}{2} \frac{(\delta m^2)^2}{\lambda} \int_{-\infty}^{+\infty} \left[1 - \cos \frac{\sqrt{\lambda}}{m} \phi_{\text{sol}}(x) \right] dx = \frac{\lambda}{8mL^2} \left(\sum \frac{1}{\tilde{\omega}} \right)^2 \quad (128)$$

where we wrote δm^2 as $\frac{\lambda}{4L} \sum \frac{1}{\tilde{\omega}}$.

We first demonstrate that the terms due to expanding the denominators $(1 + q_n^2)L - \frac{2}{m}$ cancel. Consider as an example the last term in the first line of (124). It cancels the last term in the first line of (126). To see this, write δm^2 into its original form as a sum over modes

$$\delta m^2 = \frac{\hbar \lambda}{4L} \left(\sum \frac{1}{\tilde{\omega}} \right) \quad (129)$$

One finds for the terms with L^{-2}

$$\begin{aligned} & \left(-\frac{\lambda}{32} \right) \frac{4}{L^2} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\omega^3} - \frac{1}{2} \left(-\frac{\lambda}{4L} \sum \frac{1}{\tilde{\omega}} \right) \frac{1}{mL} \sum \frac{1}{\omega^3} \right) \\ &= -\frac{\lambda}{8L^2} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right) \left(\sum \frac{1}{\omega^3} \right) \end{aligned} \quad (130)$$

Note now that $\frac{1}{L} \sum \frac{1}{\omega^3}$ is finite, while

$$\begin{aligned} \frac{1}{L} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right) &= \frac{1}{L} \left[\left(\sum_{-N}^{-1} + \sum_1^N \right) \frac{1}{\omega_n} - \sum_{-N}^N \frac{1}{\tilde{\omega}_n} \right] \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{1}{\sqrt{k^2 + m^2}} \left(\frac{\delta(k)}{L} \right) dk \rightarrow 0 \end{aligned} \quad (131)$$

is unambiguous and vanishes (the absence of the $n = 0$ contribution allows the application of the Euler-Maclaurin formula as discussed after (122)). Hence, the contributions due to expanding the denominator $(q_n^2 + 1)L - \frac{2}{m}$ cancel. Similar cancelations occur in other pairs of corresponding terms, and we shall therefore only use the terms $(q_n^2 + 1)L$ in the denominators of $E(x, q_n)$ from now on.

The sum of all contributions from the terms with η^4 , $\delta m^2 \eta^2$, and $(\delta m^2)^2$ in the topological and trivial sectors is then found to combine into differences, except for one term, namely the contribution from the first term in the third line in (124)

$$\begin{aligned} \langle H^{(2)} \rangle \Big|_{\eta^4 \text{ and } \delta m^2 \eta^2} &= \begin{array}{c} \text{Diagram 1} + \text{Diagram 2} - \left\{ \text{Diagram 3} + \text{Diagram 4} \right\} + (\blacksquare)^2 = \\ -\frac{\lambda}{32L} \left(\sum \frac{1}{\omega} \right)^2 + \frac{\lambda}{16L} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\tilde{\omega}} \right) + \left(\frac{\lambda}{32} - \frac{\lambda}{16} \right) \frac{1}{L} \left(\sum \frac{1}{\tilde{\omega}} \right)^2 \\ + \frac{\lambda}{8mL^2} \left(\sum \frac{1}{\omega} \right)^2 - \frac{\lambda}{4mL^2} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\tilde{\omega}} \right) + \frac{\lambda}{8mL^2} \left(\sum \frac{1}{\tilde{\omega}} \right)^2 \\ - \frac{\lambda m}{24L^2} \left(\sum \frac{1}{\omega} \right) \left(\sum \frac{1}{\omega^3} \right) + \frac{\lambda m}{24L^2} \left(\sum \frac{1}{\omega^3} \right) \left(\sum \frac{1}{\tilde{\omega}} \right) \\ + \frac{\lambda m^3}{40L^2} \left(\sum \frac{1}{\omega^3} \right)^2 \end{array} \\ &= -\frac{\lambda}{32L} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right)^2 + \frac{\lambda}{8mL^2} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right)^2 \\ &\quad - \frac{\lambda m}{24L^2} \left(\sum \frac{1}{\omega^3} \right) \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right) + \frac{\lambda m^3}{40L^2} \left(\sum \frac{1}{\omega^3} \right)^2 \end{aligned} \quad (132)$$

Drawn lines represent propagators in the soliton sector and dotted lines propagators in the trivial vacuum. In the intermediate expression the first column gives the contributions from the η^4 term in $H_{\text{int, sol}}^{(2)}$ while the second column gives the contribution from $\delta m^2 \eta^2$ in ΔH_{sol} . The third column contains the contributions from the vacuum sector (in the first row) and from the term $(\delta m^2)^2$ in the topological sector (in the second row). We recall that ω and $\tilde{\omega}$ denote the frequencies in the topological and trivial sectors respectively. We claim that all differences cancel. Since we already proved this for (131), we only need to discuss the sum $\frac{1}{L} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right)^2$. Each factor $\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}}$ is ambiguous but finite, see (131). Hence the extra $\frac{1}{L}$ factor ensures that also this term vanishes.

We conclude that all η^4 , η^2 and η^0 terms contribute only one term

$$\frac{\lambda m^3}{40L^2} \left(\sum \frac{1}{\omega^3} \right)^2 = \frac{\lambda}{40\pi^2 m} \quad (133)$$

This is the contribution due to the one-vertex two-loop graph in which one only retains in both propagators the deviations from the trivial space propagators. The individual other terms are ambiguous and divergent but their sum cancels.

The η^3 and $\delta m^2 \eta$ contributions with one intermediate particle

We next evaluate the contributions to the mass of the soliton which come from the η^3 term in $H_{\text{int,sol}}^{(1)}$, and the η term in ΔH_{sol} . We first take one-particle intermediate states in the sums over $|p\rangle$ in (1). Since there are no terms in the trivial sector which are odd in η , these contributions should sum up to a finite result, but we are again interested in possible ambiguities. Using

$$\begin{aligned} & \langle n | \frac{\delta m^2 m}{\sqrt{\lambda}} \int_{-\infty}^{+\infty} \frac{-4(e^{mx} - e^{-mx})}{(e^{mx} + e^{-mx})^2} \eta dx | \text{sol} \rangle \\ &= -\frac{\delta m^2 \pi}{m \sqrt{2L\lambda}} \frac{\sqrt{\omega_n}}{\cosh \frac{1}{2} \pi q_n} \end{aligned} \quad (134)$$

(see the appendix), and

$$\frac{1}{L} \sum_n \frac{1}{\cosh^2(\frac{1}{2} \pi q_n)} = \frac{2m}{\pi^2} \quad (135)$$

we find straightforwardly

$$\sum_n |\langle n | \eta \text{ term} | \text{sol} \rangle|^2 / (-\omega_n) = -\frac{(\delta m^2)^2}{\lambda m} \quad (136)$$

Next we evaluate $\langle p | \eta^3 \text{ term} | \text{sol} \rangle$ with $\langle p |$ a one-particle state. This yields a factor of 3 times an equal-time contraction at the point x . The x -integrals are given in the appendix and one finds

$$\langle p, 1 \text{ part} | \eta^3 \text{ term} | \text{sol} \rangle = \sqrt{\frac{\lambda}{2L}} \frac{\pi}{4m} \sum_p \frac{\sqrt{\omega_p}}{\cosh \frac{1}{2} \pi q_p} \left(\frac{1}{L} \sum_n \frac{1}{\omega_n} - \frac{\omega_p^2}{4\pi m^2} \right) \quad (137)$$

Then it is relatively easy to obtain

$$\begin{aligned} & \sum_p \frac{2}{-\omega_p} \text{Re} \langle \text{sol} | \eta^3 \text{ term} | p, 1 \text{ part} \rangle \langle p, 1 \text{ part} | \eta \text{ term} | \text{sol} \rangle \\ &= (\delta m^2) \left[-\frac{1}{2mL} \sum \frac{1}{\omega} - \frac{1}{6\pi m} \right] \end{aligned} \quad (138)$$

Finally we evaluate

$$\begin{aligned} & \sum_p \frac{1}{-\omega_p} |\langle p, 1 \text{ part} | \eta^3 \text{ term} | \text{sol} \rangle|^2 \\ &= -\frac{1}{\lambda m} \left(\frac{\lambda}{4L} \sum \frac{1}{\omega} \right)^2 + \frac{1}{6\pi m} \left(\frac{\lambda}{4L} \sum \frac{1}{\omega} \right) - \frac{\lambda}{120\pi^2 m} \end{aligned} \quad (139)$$

The sum of the contributions from the $\delta m^2 \eta$ and η^3 terms with only one-particle intermediate states is then

$$\begin{aligned}
& \blacksquare \text{---} \blacksquare + \blacksquare \text{---} \bigcirc + \bigcirc \text{---} \bigcirc = \\
& -\frac{1}{\lambda m}(\delta m^2)^2 + \delta m^2 \left(\frac{1}{2mL} \sum \frac{1}{\omega} - \frac{1}{6\pi m} \right) - \frac{1}{\lambda m} \left(\frac{\lambda}{4L} \sum \frac{1}{\omega} \right)^2 \\
& + \frac{1}{6\pi m} \left(\frac{\lambda}{4L} \sum \frac{1}{\omega} \right) - \frac{\lambda}{120\pi^2 m} \\
& = -\frac{\lambda}{16mL^2} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right)^2 + \frac{\lambda}{24\pi mL} \left(\sum \frac{1}{\omega} - \sum \frac{1}{\tilde{\omega}} \right) \\
& - \frac{\lambda}{120\pi^2 m}
\end{aligned} \tag{140}$$

The only nonvanishing contribution from these terms is thus due to the square of the finite part of the matrix element of the η^3 term, and the latter is again obtained by taking the deviation from the propagator in the trivial vacuum (the part proportional to $\tanh^2 mx - 1$) in the equal-time contraction of two η fields. All other matrix elements are divergent, and none of them contributes.

The contribution from three intermediate particles and the $\eta\pi\eta\pi$ term

From (111) the matrix element with 3 intermediate particles can be written as follows (after the substitution $e^{mx} = y$)

$$\begin{aligned}
& \frac{1}{2m} \int_0^\infty dy^2 \left[\frac{(y^2 - 1)^4}{(y^2 + 1)^5} - i \frac{(y^2 - 1)^3}{(y^2 + 1)^4} (q_n + q_r + q_t) \right. \\
& \left. - \frac{(y^2 - 1)^2}{(y^2 + 1)^3} (q_n q_r + q_n q_t + q_r q_t) + i \frac{y^2 - 1}{(y^2 + 1)^2} q_n q_r q_t \right] (y^2)^{\frac{i}{2}Q - \frac{1}{2}}
\end{aligned} \tag{141}$$

where $Q = q_n + q_r + q_t$. Using

$$\int_0^\infty dy^2 y^n y^{iQ-1} (y^2 + 1)^{-m} = B\left(\frac{iQ}{2} + \frac{n+1}{2}, \frac{-iQ}{2} - \frac{n+1}{2} + m\right) \tag{142}$$

where B is the beta function, one finds

$$\begin{aligned}
& \frac{1}{2m} \left[-\frac{1}{8}Q^4 + \frac{1}{4}Q^2 + \frac{1}{2}(Q^2 - 1)(q_n q_r + q_n q_t + q_r q_t) - Q(q_n q_r q_t) + \frac{3}{8} \right] \frac{\pi}{\cosh \frac{1}{2}\pi Q} \\
& = \frac{1}{2m} \left[-\frac{1}{8}(\omega_n^2 + \omega_r^2 + \omega_t^2)^2 + \frac{1}{2}(\omega_n^2 \omega_r^2 + \omega_n^2 \omega_t^2 + \omega_r^2 \omega_t^2) \right] \frac{\pi}{\cosh \frac{1}{2}\pi Q}
\end{aligned} \tag{143}$$

Squaring, one finds for the contributions to $M^{(2)}$ due to intermediate states with 3 particles

$$\begin{aligned}
& \frac{1}{6} \sum_{n,r,t} \left| \langle q_n, q_r, q_t | H_{\text{int,sol}}^{(1),I} | \text{sol} \rangle \right|^2 \frac{-1}{\omega_n + \omega_r + \omega_t} = \bigcirc \\
& = \frac{-\lambda}{2^{10}\pi 6m} \int \frac{dq_1 dq_2 dq_3}{\cosh^2 \frac{1}{2}\pi(q_1 + q_2 + q_3)} \\
& \times \frac{[(1+q_1^2)^2 + (1+q_2^2)^2 + (1+q_3^2)^2 - 2(1+q_1^2)(1+q_2^2) - 2(1+q_1^2)(1+q_3^2) - 2(1+q_2^2)(1+q_3^2)]^2}{(1+q_1^2)^{3/2}(1+q_2^2)^{3/2}(1+q_3^2)^{3/2}(\sqrt{1+q_1^2}\sqrt{1+q_2^2}\sqrt{1+q_3^2})}
\end{aligned} \tag{144}$$

The prefactor $\frac{1}{6}$ is needed since we sum over all q_n, q_r, q_t while the 3-particle states are given by $a_n^\dagger a_r^\dagger a_t^\dagger |\text{sol}\rangle$.

Next we evaluate the contribution from the $\int \eta' \pi \int \eta' \pi$ term. Using that $\eta = \sum_n (2\omega_n L)^{-1} (a_n E_n \exp -i\omega_n t + h.c.)$ while $\pi = \sum_n (2\omega_n L)^{-1} (-i\omega_n a_n E_n \exp -i\omega_n t + h.c.)$ one finds 3 contributions from the 3 possible contractions,¹⁴

$$\begin{aligned} \langle \text{sol} | H_{\text{int,sol}}^{(2),III} | \text{sol} \rangle &= \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} \\ &= \frac{\lambda}{64m^3} \sum_{n,r} \left(-C_{n,n} C_{r,r} + C_{r,-n} C_{-n,r} + \frac{\omega_r}{\omega_n} C_{n,-r} C_{-n,r} \right) \end{aligned} \quad (145)$$

where $C_{n,r} = \int (dE_n/dx) E_r dx$. Dashed lines denote π fields. Using that E_n and E_r are orthonormal, it follows that

$$C_{n,r} = imq_n \delta_{n,r} + \frac{1}{L} \frac{1}{\omega_n \omega_r} (q_n^2 - q_r^2) \frac{i\pi}{\sinh \frac{1}{2}\pi(q_n - q_r)} \quad (146)$$

Since $C_{n,n}$ is odd in n ($C_{n,n} = imq_n + 2iq_n/(L\omega_n^2)$) the contribution from $C_{n,n} C_{r,r}$ vanishes, and in the remainder the term with $imq_n \delta_{n,r}$ cancels in the contribution $\sum_{r,n} (\frac{\omega_r}{\omega_n} - 1) |C_{n,r}|^2$. This leads to

$$\begin{aligned} &\langle \text{sol} | \int \eta' \pi \int \eta' \pi \text{-term} | \text{sol} \rangle \\ &= \frac{\lambda}{2^{10}m} \int \frac{dq_1 dq_2}{[\sinh \frac{1}{2}\pi(q_1 - q_2)]^2} \left(\frac{\sqrt{1+q_1^2}}{\sqrt{1+q_2^2}} - 1 \right) \frac{(q_1^2 - q_2^2)^2}{(1+q_1^2)(1+q_2^2)} \end{aligned} \quad (147)$$

According to Verwaest [20] the sum of these two contributions is

$$- \frac{\lambda}{60\pi^2 m}. \quad (148)$$

(In ref. [19] these contributions were numerically evaluated). For us the crucial point is that both contributions are finite and hence unambiguous.

Adding (133), (140) and (148) one finds that the sum of these contributions vanishes. Hence, the only contribution to the two-loop correction of the soliton mass comes from the term $-\hbar^2/(8M_{\text{cl}}^2) \int (\phi'')^2 dx$ in (106) and it yields

$$M^{(2)} = -\frac{\lambda}{192m}. \quad (149)$$

All other contributions combine into unambiguous finite integrals which vanish due to factors $1/L$.

¹⁴ Each of these 3 contributions is proportional to $\hbar^2$ because each propagator gives a factor $\hbar$, vertices do not contribute factors of $\hbar$, and the energy denominators in (77) yield a factor $\hbar^{-1}$.

7 Conclusions

In this article two new concepts have been introduced in the theory of quantum solitons: topological boundary conditions and a physical principle which fixes UV quantum ambiguities. The main idea underlying these concepts is simple: the problem must be formulated *before* the loop expansion is performed. This means that the mass of the soliton must be defined nonperturbatively, rather than as a sum of the classical result, plus one-loop corrections, plus etc. We define this mass as the difference between the vacuum energies of the system with different boundary conditions. It follows immediately that the boundary conditions must be formulated for the full quantum field, rather than for small fluctuations.

The topological boundary conditions are, in fact, better viewed as conditions which put the system on a Moebius strip without boundaries. In the literature one usually employs periodic boundary conditions in the soliton sector for all the quantum fluctuations (but not for the classical field), or a mixture of periodic conditions for bosons and others for fermions. All these conditions may distort the system at the boundaries and introduce spurious extra energy $O(L^0)$ contributions which obscure the measurement of the mass of a soliton. We even found that most boundary conditions in the literature are not compatible with the Majorana condition for fermions in the $N = (1, 1)$ case. With topological boundary conditions there is no spurious energy introduced at the boundaries since there are no boundaries, and one obtains the genuine mass of a soliton.

In the long-standing problem of UV ambiguities of the quantum mass of a soliton, we have taken the point of view that this problem should be recognized as the well-known regularization dependence of loop calculations in quantum field theory. The action is determined only up to local counterterms, and one has to introduce renormalization conditions to fix those. With solitons, we want to compare vacuum energies in topologically distinct sectors. The new principle we have introduced is that if all mass parameters in the theory tend to zero, the topological and the trivial sector must have the same vacuum energy (in the infinite volume limit).

This condition follows immediately from a simple dimensional analysis. Indeed, if the physical meson mass m and all dimensionful couplings scale according to their dimensions as $m \rightarrow 0$, then the mass of the soliton M as a function of these parameters must also scale as m . One can take a different, but equivalent, look at the problem: After rotation to Euclidean space one finds that the soliton is a string-like interface between two phases in two-dimensional classical statistical field theory. Our principle is then related to the well-known property of the interface tension: the tension vanishes at the critical (conformal) point [17]. Note again that the formulation of this principle is entirely nonperturbative.

It is when we do the loop expansion we see that the principle leads to nontrivial consequences. This is because divergent contributions proportional to the UV cutoff Λ and independent of m may appear (and they do appear in the $N = 1$ theory with mode number cutoff and topological boundary conditions). Our principle can be used as a renormalization condition to eliminate such divergences. Adopting this principle, the calculation of the soliton mass becomes unambiguous if one first differentiates the sums over frequencies w.r.t. the mass parameter, and then integrates setting the integration constant equal to zero.

There exist other methods to compute masses in certain 1+1-dimensional models [32,

35, 36, 37, 38, 39, 21, 22]. For exactly solvable models one can determine the S-matrix exactly by assuming that the Yang-Baxter equation holds and that the S-matrix factorizes into products of two-particle S-matrices. This program has been extended to the $N = 1$ susy sine-Gordon model [21, 22] and from the result one reads off that to one loop the soliton mass is given by

$$M = M_{\text{cl}}(m) - \frac{m}{2\pi}, \quad (150)$$

where m is the physical renormalized meson mass. This is the same result as we have found with topological boundary conditions and our renormalization condition at the conformal point. From our point of view, choosing the Yang-Baxter equation (or the thermodynamic Bethe ansatz together with the quantum Lax-pairs approach — the “inverse scattering method”) amounts to a choice of regularization scheme, which, at least as far as the quantum mass of solitons is concerned, appears to be equivalent to our principle.

However, it may seem that the BPS bound $\langle H \rangle \geq |\langle Z \rangle|$ is violated at the one-loop level because of the negative sign of $M^{(1)}$, since the central charge does not receive quantum corrections at one loop [8, 14] (apart from those absorbed by the renormalization of m). We point out that it is the unrenormalized expectation value $\langle H \rangle$ of the Hamiltonian that should obey the bound, not the physical soliton mass M , which may (and does in the $N = 1$ case) differ from $\langle H \rangle$ by an m -independent counterterm. Using the mode-number cut-off regularization and topological boundary conditions, we have found that

$$\langle H \rangle_{\text{MNC}} = M_{\text{cl}}(m) + \frac{\Lambda}{4} - \frac{m}{2\pi} \quad (151)$$

where Λ is the ultraviolet cut-off. This means that the bound is observed by $\langle H \rangle$. This happens because $N = 1$ susy is not enough to eliminate the linear divergent term in $\langle H \rangle$. This term is positive in accordance with the BPS bound. The bound is not saturated, which agrees with the observation of Olive and Witten [3] that the saturation of the bound is related to “multiplet shortening”. The latter does not occur in the $N = 1$ model.

Therefore, in the $N = 1$ theory with mode number cutoff we encounter a situation where our physical principle $M|_{m=0} = 0$ leads to nontrivial consequences. The subtraction of the trivial vacuum energy as in (3) eliminates bulk volume $O(L)$ contributions (in susy theories they are absent anyway). This subtraction cannot, however, eliminate a possible $O(L^0)$ m -independent but regularization dependent $O(\Lambda)$ contribution. The required subtraction constant, or counterterm, is determined by the condition $M|_{m=0} = 0$. Therefore

$$M = (E_1 - E_0) - (E_1 - E_0)_{m=0} \quad (152)$$

is a more complete (compared to (3)) definition of the physical soliton mass. It is clear that this is exactly the definition implemented by our d/dm calculation

$$M = \int_0^m \frac{d}{dm}(E_1 - E_0). \quad (153)$$

In $N = 2$ models we do not encounter any of these issues. There it turned out that neither the soliton mass nor the central charge receive quantum corrections, hence the BPS bound remains intact and saturated. In contrast with the $D = 2$ $N = 1$ case where all susy

representations, with and without saturation of the bound, are two-dimensional, the BPS bound of the $N = 2$ models is protected by “multiplet shortening” [3].

An alternative UV regularization which one may use is the energy cutoff. In the mode cutoff regularization one truncates the *sums* over the modes. The energy cutoff amounts to first converting the sums into integrals over momenta and then truncating these *integrals*. As was shown in [14] this regularization scheme in the sine-Gordon model leads to a result in disagreement with the Dashen-Hasslacher-Neveu spectrum [39, 16]. In the supersymmetric sine-Gordon case the energy cutoff would lead to a vanishing one-loop correction (after standard renormalization of m), which is in contradiction with existing exact results [21, 22]. We examined the two-loop corrections and found that no dependence on the choice of regularization appears there. Therefore the difference between the energy cutoff and the mode cutoff is purely a one-loop effect. This suggests that, perhaps, a formulation of the theory exists where this effect can be described in terms of a quantum topological one-loop anomaly. Moreover, the one-loop correction to the mass M does not depend on the coupling constant, thus it is, in a certain sense, a geometrical effect.

Acknowledgments

We would like to thank N. Graham and R. Jaffe for discussions.

Appendix

$$\int_{-\infty}^{+\infty} \left[\frac{8}{(e^{mx} + e^{-mx})^2} \right] dx = \frac{4}{m} \quad (154)$$

$$\int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2} \right] (\tanh^2 mx - 1) dx = \frac{2}{3m} \quad (155)$$

$$\int_{-\infty}^{+\infty} \left[1 - \frac{8}{(e^{mx} + e^{-mx})^2} \right] (\tanh^2 mx - 1)^2 dx = -\frac{4}{5m} \quad (156)$$

$$\int_{-\infty}^{+\infty} \frac{e^{mx} - e^{-mx}}{(e^{mx} + e^{-mx})^2} e^{imq_n x} (\tanh mx - iq_n) dx = \frac{\pi}{4m} \frac{1 + q_n^2}{\cosh \frac{1}{2} \pi q_n} \quad (157)$$

$$\int_{-\infty}^{+\infty} \frac{e^{mx} - e^{-mx}}{(e^{mx} + e^{-mx})^4} e^{imq_n x} (\tanh mx - iq_n) dx = \frac{1}{64m} (q_n^2 + 1)^2 \frac{\pi}{\cosh \frac{1}{2} \pi q_n} \quad (158)$$

(Substitute $e^{mx} = y$, write $iq_n e^{imq_n x}$ as $\frac{1}{m} \frac{d}{dx} e^{imq_n x}$ and put $y^2 = \frac{1}{t} - 1$. Then use beta-functions and the identities $\Gamma(1+x) = x\Gamma(x)$ and $\Gamma(x)\Gamma(1-x) = \pi/\sin \pi x$).

$$\frac{1}{L} \sum_p \frac{1}{(\cosh \frac{1}{2} \pi q_p)^2} = \frac{2m}{\pi^2} \quad (159)$$

$$\frac{1}{L} \sum_p \frac{\omega_p^2}{(\cosh \frac{1}{2} \pi q_p)^2} = \frac{8m^3}{3\pi^2} \quad (\text{Use } \int_0^\infty \frac{x dx}{e^{\pi x} + 1} = \frac{1}{12}). \quad (160)$$

$$\frac{1}{L} \sum_p \frac{\omega_p^4}{(\cosh \frac{1}{2} \pi q_p)^2} = \frac{64m^5}{15\pi^2} \quad (161)$$

References

- [1] R. Rajaraman, *Solitons and instantons* (North-Holland, Amsterdam, 1982)
- [2] N. Seiberg, E. Witten, Nucl. Phys. B431 (1994) 484; *ibid.* B426 (1994) 19; Erratum *ibid.* B430 (1994) 485.
- [3] D. Olive and E. Witten, Phys. Lett. 78B (1978) 97
- [4] A. d’Adda and P. di Vecchia, Phys. Lett. 73B (1978) 162; A. d’Adda, R. Horsley and P. di Vecchia, Phys. Lett. 76B (1978) 298; R. Horsley, Nucl. Phys. B151 (1979) 399
- [5] J. F. Schonfeld, Nucl. Phys. B161 (1979) 125
- [6] S. Rouhani, Nucl. Phys. B182 (1981) 462.
- [7] R.K.Kaul and R. Rajaraman, Phys. Lett. B131 (1983) 357
- [8] C. Imbimbo and S. Mukhi, Nucl. Phys. B247 (1984) 471
- [9] H. Yamagishi, Phys. Lett. 147B (1984) 425
- [10] A. K. Chatterjee and P. Majumdar, Phys. Rev. D30 (1984) 844; Phys. Lett. 159B (1985) 37
- [11] A. Uchiyama, Nucl. Phys. B244 (1984) 57
- [12] A. Uchiyama, Progr. Theor. Phys. 75 (1986) 1214
- [13] A. Uchiyama, Nucl. Phys. B278 (1986) 121
- [14] A. Rebhan and P. van Nieuwenhuizen, Nucl. Phys. B508 (1997) 449
- [15] S. Coleman, Phys. Rev. D11 (1975) 2088
- [16] R. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. D10 (1974) 4114 and 4130; Phys Rev D 12 (1975) 2443
- [17] D. Jasnow, in *Phase Transitions and Critical Phenomena*, ed. C. Domb and J.L. Lebowitz, Academic Press, London 1986, vol. 10
- [18] E. Farhi, N. Graham, P.E. Haagensen and R.L. Jaffe, e-print hep-th/9802015
- [19] H.J. De Vega, Nucl. Phys. B115 (1976) 411
- [20] J. Verwaest, Nucl. Phys. B123 (1977) 100
- [21] K. Schoutens, Nucl. Phys. B344 (1990) 665
- [22] C. Ahn, D. Bernard and A. LeClair, Nucl. Phys. B246 (1990) 409; C. Ahn, Nucl. Phys. B354 (1991) 57

- [23] C. Korthals Altes, A. Michels, M. Stephanov, M. Teper, Phys. Rev. D55 (1997) 1047
- [24] P. Morse and H. Feschbach, *Methods in mathematical physics*, McGraw-Hill Book Co., New York, 1953
- [25] J-L. Gervais and A. Jevicki, Phys. Rev. D11 (1975) 2943
- [26] J-L. Gervais and A. Jevicki and B. Sakita, Phys. Rev. D12 (1975) 1038
- [27] J-L. Gervais and B. Sakita, Phys Rev D16 (1977) 3507
- [28] A. Jevicki, Nucl Phys B117 (1976) 365
- [29] N. Christ and T.D. Lee, Phys. Rev. D12 (1975) 1606
- [30] J. Goldstone and R. Jackiw, Phys. Rev. D11 (1975) 1486
- [31] L. Jacobs, Phys. Rev. D13 (1976) 2278
- [32] R. Dashen, B. Hasslacher and A. Neveu, Phys. Rev. D11 (1975) 3424
- [33] E. Tomboulis, Phys. Rev. D12 (1975) 1678
- [34] J-L. Gervais and A. Jevicki, Nucl. Phys. B110 (1976) 93 and 113
- [35] A.M. Polyakov, Pisma Zh. E. T. F. 20 (1975) 430
- [36] J.D. Johnson, S. Krinsky, B.M. McCoy, Phys. Rev. A8 (1973) 2526.
- [37] I.Ya. Aref'eva and V.E. Korepin, JETP Lett. 20 (1974) 312
- [38] L.D. Faddeev and V.E. Korepin, Phys. Rep. 42 (1978) 1
- [39] L.D. Faddeev and V.E. Korepin, Theor. Math. Phys. 25 (1975) 147